

# **Bizmut a jeho alteračné produkty: akcesorické minerály granitového pegmatitu berylového typu na lokalite Bratislava - Švábsky vrch (Malé Karpaty, Slovensko)**

**Native bismuth and its alteration products: accessory minerals of beryl type granitic pegmatite from Bratislava, Švábsky Hill (Malé Karpaty Mts., Slovakia)**

MARTIN ŠTEVKO\*, PAVEL UHER A DANIEL OZDÍN

*Katedra mineralogie a petrologie, Prírodovedecká fakulta, Univerzita Komenského v Bratislave, Mlynská dolina 4,  
842 15 Bratislava, Slovenská republika; \*e-mail: stevko@fns.uniba.sk*

ŠTEVKO M., UHER P., OZDÍN D. (2012): Bizmut a jeho alteračné produkty: akcesorické minerály granitového pegmatitu berylového typu na lokalite Bratislava - Švábsky vrch (Malé Karpaty, Slovensko). *Bull. mineral.-petrolog. Odd. Nár. Muz. (Praha) 20, 2, 226-229. ISSN 1211-0329.*

## **Abstract**

Native bismuth forms rare aggregates (up to 1 mm across) in quartz core of the beryl-type granitic pegmatite from Bratislava, Švábsky Hill (Malé Karpaty Mts., Slovakia). The pegmatite dike (<1 m thickness) cuts parental Hercynian S-type granites of the Bratislava massif. The bismuth shows homogeneous, nearly end-member composition (only 0.2 wt.% Sb). It is partly replaced by bismite and bismutite along the rims. Native bismuth precipitated during final hydrothermal stage of the pegmatite crystallization at temperature 250 - 260 °C, whereas supergene bismite and bismutite are products of late oxidation and weathering processes, or low-temperature alteration of the bismuth.

**Key words:** native bismuth, bismite, bismutite, chemical composition, granitic pegmatites, Bratislava, Malé Karpaty Mts., Slovakia