

Historie dolování, ložisková geologie a mineralogie některých lokalit v novoknínské zlatonosné oblasti (střední Čechy)

Mining history, economic geology and mineralogy of some localities in the Nový Knín gold-bearing area (central Bohemia, Czech Republic)

JIŘÍ LITOCHLEB¹⁾, JIŘÍ ŠEJKORA¹⁾, VLADIMÍR ŠREIN²⁾ A BLANKA ŠREINOVÁ¹⁾

¹⁾ Mineralogicko-petrologické oddělení, Národní muzeum, Cirkusová 1740, 193 00 Praha 9 - Horní Počernice

²⁾ Česká geologická služba, Klárov 3, 118 21 Praha 1

LITOCHLEB J., ŠEJKORA J., ŠREIN V., ŠREINOVÁ B. (2011): Historie dolování, ložisková geologie a mineralogie některých lokalit v novoknínské zlatonosné oblasti (střední Čechy). - Bull. mineral.-petrolog. Odd. Nár. Muz. (Praha) 19/2, 101-129. ISSN 1211-0329.

Abstract

The Nový Knín area represents besides the Jílové gold-bearing district the second very significant gold-bearing region in the central Bohemia (Czech Republic). This paper summarizes hitherto knowledge about mining history, economic geology and mineralogy of some localities in the Nový Knín gold-bearing area and presents the results of a new mineralogical research. Gold mineralization in the Nový Knín, Kozí Hory and Libčice gold-bearing districts is represented by the veins or stockworks in sedimentary and volcanosedimentary Neoproterozoic rocks of the Barrandian block (bohemicum). The primary gold-bearing occurrences are often accompanied by alluvial and deluvial gold-bearing placers with visible gold.

In this ore area, the following types of mineralization occurs: *classic quartz-base-metals* (arsenopyrite, Fe-rich sphalerite, galena, chalcopyrite, pyrite), *quartz-Au-Ag-base-metals* (arsenopyrite, Fe-rich sphalerite, löllingite, galena, chalcopyrite, Fe and Sb dominant freibergite, hessite, acanthite, miargyrite, Au-Ag phases - gold (fineness 0.340 - 0.515) to Au-rich silver and accessory sulphosalts (Pb-Cu-Bi,Sb-S and Pb-Ag-Sb-S), *quartz-gold-bearing* (arsenopyrite, pyrite, pyrhotite, chalcopyrite, locally molybdenite, gold (fineness >0.860), Bi-Te±S phases - tsumoite, pilsenite, sulphotsumoite, joséite B, tetradyomite) and *antimony mineralization* (gudmundite, berthierite in quartz or calcite veins or disseminated in rocks). The dominant primary quartz-gold-bearing mineralization were formed in course Variscan hydrothermal, locally in conditions contact metamorphism processes (the Libčice gold deposit).

Key words: mining history, geology, mineralogy, paragenesis, gold, tellurides, sulphotellurides, freibergite, base-metal sulphides, chemical composition, the Nový Knín gold-bearing area, central Bohemia