



EARLY MIOCENE BIRDS OF SKYŘICE, CZECH REPUBLIC

Jiří Mlíkovský

Institute of Geology and Paleontology, Charles University, Albertov 6, 128 43 Praha 2, Czech Republic

Received January 19, 2000

Accepted May 10, 2000

A b s t r a c t: The early Miocene locality Skyřice yielded remains of 3 species from 3 different families of birds, incl. Podicipedidae, Anatidae and Phasianidae. The record of *Podiceps* is the oldest for the family Podicipedidae in the Old World so far.

■ Aves, Miocene, Czech Republic.

INTRODUCTION

Early Miocene lacustrine deposits at Skyřice near Most in NW Bohemia were discovered during coal mining activities in 1902 (Schlosser et Hibsich 1902, Hibsich 1910). In addition to ostracodes and freshwater molluscs (Schlosser et Hibsich 1902), the locality yielded a limited amount of vertebrate remains, including those of fishes (Laube 1903), amphibians (Laube 1903), reptiles (Schlosser et Hibsich 1902, Laube 1910), and large mammals (Schlosser et Hibsich 1902, Schlosser 1910).

The birds from Skyřice were originally mentioned by Laube (1910). They are limited to four bone fragments, all of which were identified as those of *Anas blanchardi* MILNE-EDWARDS by Laube (1910). A restudy of three surviving bones showed that three different families of birds are represented here (see below).

The age of the locality has never been identified in terms of MN-zones (sensu Mein 1990), because no micromammals were found. However, all relevant fossiliferous deposits in the vicinity of Most are MN-3 in age (Fejfar et Kvaček 1993), so that the same age can be assumed for the Skyřice locality as well.

The remains of birds from Skyřice are currently deposited in the author's collection in Praha (MP). The stratigraphy of the Neogene follows Mein (1990).

SYSTEMATIC LIST

Order Podicipediformes FÜRBRINGER, 1888

Family Podicipedidae BONAPARTE, 1831

Genus *Podiceps* LATHAM, 1787

***Podiceps walteri* (ŠVEC, 1982)**

Miobaptus walteri ŠVEC, 1982: 246, fig. 1, pl. 2: fig. 2.

Podiceps walteri (ŠVEC): Mlíkovský this paper (new combination)

Material: distal part of left tibiotarsus; MP, uncatalogued. Figured by Laube 1910, pl. 58, fig. 4a-d.

Measurements: distal width = 5.3 mm, distal depth = 5.0 mm

Comments: This species was originally described from the early Miocene (MN 4) of Dolnice (Švec 1982, 1984) in the genus *Miobaptus* ŠVEC, 1982. Subsequent research showed that it is inseparable from the modern genus *Podiceps* LATHAM, 1787 (Olson 1995, Mlíkovský unpub. results).

The record of *Podiceps walteri* from Skyřice is the oldest for the family in the Old World. The only older record is *Podiceps oligocaenus* SHUFELDT, 1915 from the Oligocene (?) of Willow Creek in

Oregon (Shufeldt 1915, Wetmore 1937, Olson 1985). The latter locality was later believed to be early Miocene in age (Brodkorb 1963), but Becker (1987) did not include it in his list of the Neogene localities of North America, apparently assigning it again to the Oligocene. In Europe, the next oldest records are the undetermined Podicipedidae from the middle Miocene (MN 7) of Steinheim in Germany (Heizmann et Hesse 1995), *Podiceps sociata* (NAVÁS, 1922) from the late Miocene (MN 9-10) of Libros in Spain (Olson 1995), and similarly aged *Podiceps miocaenicus* KESSLER, 1984 from Chişinău in Moldova (Kessler 1984), and *Podiceps* sp. from Sokolov in Ukraine (Vojinstvens'kyj 1967).

Order Anseriformes LINNAEUS, 1758

Family Anatidae LEACH, 1820

Genus *Mionetta* LIVEZEY et MARTIN, 1988

Mionetta natator (MILNE-EDWARDS, 1867)

Anas natator MILNE-EDWARDS, 1867: 148, pl. 25: figs. 14-18, 21-22.

Querquedula natator (MILNE-EDWARDS): Brodkorb 1964: 223 (new combination)

Dendrochen natator (MILNE-EDWARDS): Cheneval 1983: 93 (new combination)

Mionetta natator (MILNE-EDWARDS): Livezey et Martin 1988: 208 (new combination)

Material: carpometacarpus dex.; MP, uncatalogued. Figured by Laube 1910, pl. 58, fig. 3.

Measurements: greatest length = 36.6 mm.

Comments: This species is known also from the early Miocene (MN 2) of Saint-Gérard-le-Puy in France (Milne-Edwards 1867, Cheneval 1983, 1984a), and the early Miocene (MN 4) of Dolnice in the Czech Republic (Mlíkovský, unpub. results).

Order Galliformes LINNAEUS, 1758

Family Phasianidae HORSFIELD, 1821

Genus *Palaeortyx* MILNE-EDWARDS, 1869

Palaeortyx longipes MILNE-EDWARDS, 1869

Palaeortyx longipes MILNE-EDWARDS, 1869: 245.

Material: shaft of right humerus; MP, uncatalogued. Figured by Laube 1910, pl. 58, fig. 1a-b.

Measurements: width and depth of the midshaft = 6.3 × 5.4 mm.

Remarks: Previous early Miocene record of this species was limited to Saint-Gérard-le-Puy (MN 2a) in France (see Mlíkovský 2000), Wintershof-West (MN 3) in Germany (Ballmann 1969), and Dolnice (MN 4) in the Czech Republic (Mlíkovský, unpub. results).

DISCUSSION

Skyřice is one of the west Bohemian localities (Mlíkovský 1992, 1996a, 1998), which yielded a typical Orleanian avifauna (Mlíkovský 1996b).

The record of *Miobaptus walteri* is important from the biochronological point of view, because it represents the oldest record of the family Podicipedidae in the Old World. So far, the oldest record came from MN-zone 4b (Švec 1982, 1984, Mlíkovský 1996b).

ACKNOWLEDGMENTS

During the preparation of this study, comparative collections of recent bird skeletons were used in the Paleontological Institute of the Russian Academy of Sciences in Moscow, and in the National Museum of Natural History in Washington, D.C. I was allowed to use these collections by E. N. Kuročkin and

A. A. Karchu (Moscow), and S. L. Olson (Washington), respectively. I thank them all. My work in Washington, D.C., was sponsored by the Smithsonian Institution.

REFERENCES

- Ballmann, P. (1969): Die Vögel aus der altburdigalen Spaltenfüllung von Wintershof (West) bei Eichstätt in Bayern. – *Zitteliana* 1: 5-60.
- Becker, J. J. (1987): Neogene avian localities of North America. – Smithsonian Institution Press, Washington, 171 pp.
- Brodkorb, P. (1963): Catalogue of fossil birds: Part 1 (Archaeopterygiformes through Ardeiformes). – Bulletin of the Florida State Museum, Biological Sciences 7: 179-293.
- Brodkorb, P. (1964): Catalogue of fossil birds: Part 2 (Anseriformes through Galliformes). – Bulletin of the Florida State Museum, Biological Sciences 8: 195-355.
- Brodkorb, P. (1967): Catalogue of fossil birds: Part 3 (Ralliformes, Ichthyornithiformes, Charadriiformes). – Bulletin of the Florida State Museum, Biological Sciences 11: 99-220.
- Cheneval, J. (1983): Les Anatidae (Aves, Anseriformes) du gisement aquitainien de Saint-Gérard-le-Puy. – *In*: Buffeteaut, E., Mazin, J. M., Salmon, E. (eds.): Actes du symposium paléontologique Georges Cuvier. – Ville de Montbéliard, Montbéliard, pp. 85-98.
- Cheneval, J. (1984a): Les oiseaux aquatiques (Gaviiformes à Anseriformes) du gisement aquitainien de Saint-Gérard-le-Puy: révision systématique. – *Palaeovertebrata* 14: 33-115.
- Fejfar, O., Kvaček, Z. (1993): Tertiary basins in Northwest Bohemia. – Univerzita Karlova & Česká geologická společnost, Praha, 35 pp.
- Heizmann, E. P. J., Hesse, A. (1995): Die mittelmiozänen Vogel- und Säugetierreste des Nördlinger Ries (MN 6) und des Steinheimer Beckens (MN 7) - ein Vergleich. – *In*: Peters, D. S. (ed.): Acta palaeornithologica. Courier Forschungsinstitut Senckenberg 181: 171-185.
- Hibsch, J. E. (1910): Wichtige Funde von Wirbeltierresten im Mariannen-Schacht bei Brůx. – *Kohleninterezzent* 24: 1-3.
- Kessler, E. (1984): Noi contribuții privind studiul avifaunei din Paratethys [New contributions to the study of the Paratethys avifauna.] – *Crisia* 14: 521-532. (In Romanian).
- Laube, G. C. (1903): Batrachier- und Fischreste aus der Braunkohle von Skiritz bei Brůx. – *Sitzungsberichte des Deutschen Naturwissenschaftlich-Medizinischen Vereines für Böhmen „Lotos“ in Prag* 51: 106-114.
- Laube, G. C. (1910): Vogel- und Reptilienreste aus der Braunkohle von Skiritz bei Brůx. – *Lotos* 58: 115-127.
- Livezey, B. C., Martin, L. D. (1988): The systematic position of the Miocene anatid *Anas*[?] *blanchardi* Milne-Edwards. – *Journal of Vertebrate Paleontology* 8: 196-211.
- Mein, P. (1990): Updating of MN zones. – *In*: Lindsay, E. H., Fahlbusch, V., Mein, P. (eds.): European Neogene mammal chronology. Plenum Press, New York, pp. 73-90.
- Milne-Edwards, A., (1863): Mémoire sur la distribution géologique des oiseaux fossiles et description de quelques espèces nouvelles. – *Annales des Sciences Naturelles (Paris)* 20: 132-176.
- Milne-Edwards, A. (1867-1868): Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France. Vol. 1. – Victor Masson et Fils, Paris, 475 pp + Atlas.
- Milne-Edwards, A. (1869-1871): Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France. Vol. 2. – G. Masson, Paris, 632 pp + Atlas.
- Mlíkovský, J. (1992): The present state of knowledge of the Tertiary birds of Central Europe. – *In*: Campbell, K. E. (ed.): Studies in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County, Science Series 36: 433-458.
- Mlíkovský, J. (1996a): Tertiary avian localities of the Czech Republic. – *In*: Mlíkovský, J. (ed.): Tertiary avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 551-557.
- Mlíkovský, J. (1996b): Tertiary avian faunas of Europe. – *In*: Mlíkovský, J. (ed.): Tertiary avian localities of Europe. *Acta Universitatis Carolinae, Geologica* 39: 777-818.
- Mlíkovský, J. (1998): Early Miocene birds of Břešťany, Czech Republic. – *Časopis Národního Muzea, Řada přírodovědná* 167: 103-109.
- Mlíkovský, J. (2000): Early Miocene quails (Aves: Phasianidae) of Saint-Gérard-le-Puy, France. – *Časopis Národního Muzea, Řada přírodovědná* 169: 91 – 96.
- Navás, L., (1922): Algunos fósiles de Libros (Teruel). – *Boletín de la Sociedad Ibérica de Ciencias Naturales* 21(4): 172-174.
- Olson, S. L. (1985): The fossil record of birds. – *In*: Farner, D. S., King, J. R., Parkes, K. C. (eds.): Avian biology. Vol. 8. New York: Academic Press, pp. 79-238.

- Olson, S. L. (1995): *Thiornis sociata* Navás, a nearly complete Miocene grebe (Aves: Podicipedidae). – In: Peters, D. S. (ed.): Acta palaeornithologica. – Courier Forschungsinstitut Senckenberg 181: 131-140.
- Schlosser, M. (1910): Über fossile Wirbeltierreste aus dem Brüxer Braunkohlenbecken. – Lotos 58: 229-246.
- Schlosser, M., Hibsich, J. E. (1902): Eine untermiozäne Fauna aus dem Teplitzer Braunkohlenbecken. – Sitzungsberichte der Kaiserlich-Königlichen Akademie der Wissenschaften (Wien), Mathematisch-Naturwissenschaftliche Klasse 111: 1123-1152.
- Shufeldt, R. W. (1915): Fossil birds in the Marsh collection of Yale University. – Transactions of the Connecticut Academy of Sciences 19: 1-110.
- Švec, P. (1982): Two new species of diving birds from the Lower Miocene of Czechoslovakia. – Časopis pro Mineralogii a Geologii 27: 243-260.
- Švec, P. (1984): Further finds of grebe *Miobaptus walteri* in the Miocene of Bohemia. – Časopis pro Mineralogii a Geologii 29: 167-170.
- Vojinstvens'kyj, M. A. (1967): Iskopaemaja ornitofauna Ukrainy [Fossil avifauna of Ukraine.] – Prirodna-ja Obstanovka i Fauna Prošlogo 3: 3-76. (In Russian).
- Wetmore, A. (1937): A record of the fossil grebe, *Colymbus parvus*, from the Pliocene of California, with remarks on other American fossils of this family. – Proceedings of the California Academy of Sciences (4) 23: 195-201.

RANĚ MIOCÉNNÍ PTÁCI ZE SKYŘIC, ČESKÁ REPUBLIKA

Na raně miocénní lokalitě Skyřice u Mostu byly nalezeny kosterní zbytky tři druhů ptáků ze tří čeledí: Podicipedidae, Anatidae a Phasianidae. Nález potápky druhu *Podiceps walteri* je nejstarším dokladem čeledi Podicipedidae ve Starém Světě.