

ORDOVICIAN TERGOMYIA AND ISOSTROPHIC GASTROPODA (MOLLUSCA) OF BOHEMIA: TYPES AND REFERRED SPECIMENS IN THE COLLECTIONS OF THE NATIONAL MUSEUM, PRAGUE, CZECH REPUBLIC

RADVAN J. HORNÝ

Department of Palaeontology, National Museum, 115 79 Praha 1, Czech Republic



Horný, R. J. (2002): Ordovician Tergomyia and isostrophic Gastropoda (Mollusca) of Bohemia: Types and referred specimens in the collections of the National Museum, Prague, Czech Republic. – Acta Mus. Nat. Pragae, Ser. B, Hist. Nat., 57 [2001] (3-4): 69-102, Praha. ISSN 0036-5343.

A b s t r a c t. A review and illustrated catalogue of Bohemian Ordovician Tergomyia (Monoplacophora of previous usage) and isostrophic and patelliform Gastropoda from the Barrandian Area, Czech Republic, deposited in the collections of the National Museum, Prague. Besides descriptions, the most important data concerning the synonymy, stratigraphic range, and category of types of all valid species, supplemented by photographs, are given.

■ Mollusca, Tergomyia, isostrophic Gastropoda, Ordovician, biostratigraphy, palaeoecology, communities, mode of life, catalogue, types, referred specimens, Bohemia, Barrandian Area, Czech Republic

Received September 16, 2001

Introduction

Lower Palaeozoic Tergomyia and isostrophic Gastropoda are common molluscs, distributed in shallow-water marine deposits all over the world. During the last decade, several workers contributed to the better knowledge of these difficult and inconsistent groups, e.g. Peel (1991, preceded and followed by a series of special papers), Wahlman (1992), Webers, Pojeta, and Yochelson (1992), Geyer (1994), and Ebbestad (1999; a collection of papers summarized in a doctoral dissertation). Horný continued his studies of 1963 and since 1990 produced a number of papers.

This paper is a brief review of all to date recognized Bohemian Ordovician species of Tergomyia (Monoplacophora of previous usage) and isostrophic Gastropoda, descriptions of which are scattered among various papers starting with J. Perner (1903). Because of present importance of these groups of molluscs in evolutionary interpretations and imperative need of their study by foreign students, the catalogue is accompanied with photographs, notes, and brief descriptions of all valid taxa. The reader will find necessary data concerning biostratigraphy of the Ordovician in the Barrandian Area, as well as references of literature. All specimens have been supplemented with catalogue numbers, and various errors in numbers and indication of paralectotypes in old papers (e.g. Horný 1963a, b) have been corrected.

The catalogue is a component of a project in which the National Museum, Museum of Natural History – Department of Palaeontology has been progressively publishing catalogues of fossil flora and fauna, deposited in its collections. The author hopes to prepare a similar effort on post-Ordovician material.

The paper does not concern the intricate situation around the systematics of relevant univalve Lower Palaeozoic molluscs but largely follows the scheme published by Peel (1991).

Representatives of the Classes Tergomyia and Gastropoda constitute an important part of the Czech Ordovician benthic and epiplanktic fauna. The majority of them are interpreted as having lived in a shallow photic zone where they inhabited

either silty to sandy substrates in higher energy conditions, or quiet environments of clayey sedimentation. Specific conditions offered local bays characterized by biochemical deposition with limited influx of terrigenous material, where various types of ferrolites originated. Important is the absence of limestone sedimentation, connected with palaeoclimatic conditions in the Peri-Gondwanan province.

The gastropod and tergomyan communities contained deposit and filter feeders (*Sinuitopsis* PERNER, 1903), common epibenthic raspers, grazers, and scavengers, suprabenthic to epiplanktic species adapted to life on algae (*Tropidodiscus* MEEK et WORTHEN, 1866), and even active predators (*Sinuites* KOKEN, 1896). Some species were adapted to semifaunal mode of life (*Sinuites*). Juveniles of many species dwelled in well-aerated zone in carpets of floating algae whilst some species adapted this pseudoplanktonic mode of life even in maturity. A large diversification and enormous populations of certain species in some periods testify to rich trophic conditions, connected with influx of mineral nutrients from the nearby land and, consequently, with the development of algae.

Most finds of tergomyans and gastropods come from taphocoenoses, containing rich but usually fragmentary fauna of trilobites, brachiopods, bivalves, ostracods, carpoids, cystoids, hyoliths, conulariids, graptolites, and other groups. Rarely occur shells preserved in presumed living position, like almost sedentary limpet-like *Pygmaeconus* HORNÝ, 1961 or the semifaunal *Sinuites*. Ferrolites, locally with accumulations of shells (e.g., *Carcassonnella*), offer an extraordinary mode of preservation. Best preserved shells come from weathered, originally carbonate concretions, occurring in various strata.

The correlation with extra-Barrandian tergomyan and gastropod fauna is difficult; it is mainly caused by a limited modern study of these complex groups in the majority of neig-

hbouring regions (e.g. Sardinia, France, Spain, Portugal, Great Britain, or Russia). As a whole, the Barrandian gastropod and tergomyan fauna belongs to the Peri-Gondwanan province, with episodic influence of rare thermophilic immigrants from



Joachim Barrande, 1799-1883

Those who contributed to the knowledge of Ordovician gastropods and gastropod-like molluscs, their systematics, nomenclature, functional morphology, biology, and biostratigraphy in the post-Pernerian period were in particular R. Hor-



Jaroslav Perner, 1869-1947

the Balto-Scandinavian region, particularly in the uppermost Ordovician (Havlíček 1989, Havlíček, Vaněk, and Fatka 1994, Horný 1997g).

Joachim Barrande knew the majority of species. His incomplete manuscript and plates of the monograph *Système silurien du centre de la Bohême*, Part IV, *Gastropoda*, was finished and edited by J. Perner in three parts, issued in 1903, 1907, and 1911. The majority of old finds come from Barrande's localities, nowadays mostly inaccessible. Nevertheless, their stratigraphy is well known thanks to Barrande's descriptions, lithology, and the modern, detailed geological maps. Since Barrande's and Perner's time, the collections of gastropods and gastropod-like molluscs noticeably increased in connection with the progress of palaeontological, biostratigraphic, and geological research of the Barrandian Area. Many specimens were found completing the already described species, and numerous new species were also discovered. The modern collectors picked up all fossils that then gave a better idea of abundance and environment and did not simply concentrate on "collecting trilobites". These were namely K. Holub, F. Hanuš, C. Klouček, R. Růžička, B. Bouček, V. Plas and B. Plasová, J. Šulc, J. Putzker, J. Kraft, L. Marek, M. Šnajdr, V. Havlíček, J. Vaněk, P. Kraft, M. Mergl, R. Horný, and others. The material has been gradually housed in three main depositories – the National Museum, Prague, Museum of Dr B. Horák, Rokycany, and Czech Geological Survey, Prague. The present Catalogue concerns the material deposited in the National Museum, Prague.

ný (since 1961), L. Marek (1963), V. Havlíček and J. Vaněk (1966), and J. Frýda (since 1988). Of the modern foreign authors, James B. Knight (1941) published and re-figured the type species of genera based on Czech Lower Palaeozoic material, and John S. Peel discussed and figured two species (1974, 1991).

Acknowledgements

The paper has been prepared since 1994. I am grateful to all, who helped me with the study and preparation of the text and illustrations, as well as with important and fruitful discussions. These were John S. Peel (Uppsala University, Sweden), Jan Ove R. Ebbestad (Uppsala University, Sweden), Ivo Chlupáč (Charles University, Prague), †Vladimír Havlíček, Praha, Jaroslav Kraft, Museum of Western Bohemia, Plzeň, †Ladislav Marek, Praha, Michal Mergl, University of Western Bohemia, Plzeň, Miroslava Šandová and Martina Korandová, Museum of Dr B. Horák, Rokycany, and Petr Budil, Czech Geological Survey, Praha. Ivo Chlupáč and Ellis L. Yochelson kindly read the manuscript. I greatly appreciate many-sided help given by members of the Department of Palaeontology, National Museum, Prague: Vojtěch Turek, Rudolf Prokop, Jiří Kvaček, Kamil Zágoršek, Renata Pátová, Nadežda Zdobnická, and Ivan Kolobaba. The research had been partly supported by the Grant Agency of the Czech Republic, Grant No. 205/94/

0759 (1994–1996), and finished within the framework of the project MK0CEZ99F0201.

Systematic palaeontology

The following part, divided into two paragraphs (Tergomya; symmetrical and isostrophic Gastropoda), comprises short descriptions of all published species found in the Ordovician of Bohemia. Within the higher taxa, all subspecies are arranged in stratigraphic order. The brief synonymy is reduced to nomenclaturally important papers, as a detailed synonymy, containing all necessary data, is given in full in each item within the catalogue. Only the main or most important localities are given.

TERGOMYIA

The Class Tergomyia, previously called Monoplacophora (abandoned by Peel 1991) is represented by two orders in the Czech Ordovician: Tryblidiida and Cyrtoneillida. Only one genus, *Pentalina* HORNÝ, 1961, belongs to the tryblidiids, which are relatively common in the Silurian. The cyrtonellids are quite common, being represented by six genera and at least by ten species. With an exception of abundant *Sinuitopsis* PERNER, 1903 in the Berouian and in some strata common *Cyrtodiscus* PERNER, 1903, all other genera (*Sinuitopsis* HORNÝ, 1997, *Sarkanella* HORNÝ, 1997, *Peelerophon* YOCHELSON, 1982, and *Carcassonnella* HORNÝ et PEEL, 1996) are rare.

A favourable preservation in more or less weathered concretions, often silty and decalcified, enabled a detailed study of muscle insertions at the surface of internal moulds of *Sinuitopsis* and *Cyrtodiscus*. The functional morphology analysis brought important data, supporting the conception of W. Wenz (1940) about exogastric character of these shells, originally based on *Cyrtoneilla* HALL, 1879 and latter recognized in *Cyrtolites* CONRAD, 1838, *Yochelsonellis* HORNÝ, 1966, and others. The retractor muscle insertions are arranged in a circle running across the shell about half a whorl back from the aperture; the largest scars constitute the dorsal pair, whilst the weakest scars are in ventral position. During the growth of the shell, the muscle insertions do not move continuously as in gastropods, but periodically saltate. Besides that, the exogastric character of cyrtonellid shells is also supported by the presence of a pair of inhalant sinuses and another pair of shallow sinuses, interpreted as tentacular and therefore anterior (Horný 1991b).

Carcassonnella, which occurs in the Middle Ordovician, is closely related with the Arenigian *C. courtoissolei* (YOCHELSON, 1982) and thus connects the Mediterranean lineage. The related *Sarkanella* appears in the Middle Ordovician and can be followed up to top of the Upper Ordovician. *Peelerophon* occurs rarely in the highest Arenigian.

Of the extra-Barrandian tergomyans, types and referred specimens of Horný's Morocco paper (1997g) are deposited in the National Museum, Prague (genera *Sinuitopsis*, *Cyrtodiscus*, *Quasisinuites* HORNÝ, 1997, *Carcassonnella*, *Tachillanella* HORNÝ, 1997, and *Thoralispira* HORNÝ et VIZCAÍNO, 1995).

Order Tryblidiida LEMCHE, 1957

Family Tryblidiidae PILSBRY in ZITTEL-EASTMAN, 1899

Pentalina HORNÝ, 1961

Type species: *Pentalina prantli* HORNÝ, 1961. Middle Ordovician, Bohemia.

Pentalina prantli HORNÝ, 1961

Pl. 1, fig. 1

- 1938 *Tryblidium* sp.; Říha, Příspěvek k poznání...
1957 Tryblidiid sp.; Horný, Tryblidiinae...
1961 *Pentalina prantli* n. sp.; Horný, New genera...
1963a *Pentalina prantli* HORNÝ, 1961; Horný, Lower Palaeozoic Monoplacophora...

A rare species, so far found in two specimens. A small, up to 6.6 mm long, spoon-shaped shell with five pairs of retractor muscle scars in dorsal position. Outer shell sculpture unknown.

Stratigraphy, localities. Middle Ordovician, Dobrotivian, Dobrotivá Formation: Malé Přílepy. ? Llanvirnian, Šárka Formation: Praha – Vokovice.

Order Cyrtoneillida HORNÝ, 1963

Family Cyrtolitidae S. A. MILLER, 1889

Sinuitopsis PERNER, 1903

Type species: *Sinuitopsis neglecta* Perner, 1903. Upper Ordovician, Bohemia.

Sinuitopsis evoluta (PERNER, 1903)

Pl. 1, fig. 2

- 1903 *Sinuites sowerbyi* var. *evoluta* mihi; Perner, Système silurien, IV, Gastéropodes, 1
1963b *Sinuitopsis neglecta* PERNER; Horný: Lower Paleozoic Belle-rophontina...
1997c *Sinuitopsis?*; Horný: New, rare, ...

A rare species. According to gross morphology of the original, unique internal mould, it may be classified as *Sinuitopsis*. Two similar internal moulds come from Úvaly. The find from the Dobrotivá Formation with patches of outer shell sculpture, briefly described and figured by Horný (1997c), may also belong to this taxon. Length up to about 30 mm.

Stratigraphy and locality. Middle Ordovician, Llanvirnian, Šárka F.: Osek, Úvaly. ? Dobrotivian, Dobrotivá F.: Praha – Šárka.

Sinuitopsis neglecta

BARRANDE in PERNER, 1903

Pl. 1, figs 3, 4

- 1903 *Sinuitopsis neglecta* BARR. sp.; Perner: Système silurien, IV, Gastéropodes, 1
1903 *Sinuitopsis neglecta* BARR. var. *transgrediens* PERNER; Perner: Système silurien, IV, Gastéropodes, 1

- 1903 *Tennodiscus ferrigena*; Perner, Système silurien, IV, Gastéropodes, 1
 1903 *Tennodiscus platynotus* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
 1903 *Tennodiscus bicarinatus* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
 1903 *Tennodiscus incertus* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1941 *Sinuitopsis neglecta* PERNER; Knight, Paleozoic gastropod genotypes
 1963b *Sinuitopsis neglecta* PERNER; Horný, Lower Palaeozoic Bellerophontina...
 1991b *Sinuitopsis neglecta* PERNER; Horný, Shell morphology...

A characteristic and common cyrtonellid tergomyan, present in various sediments - clayey shales, carbonate siltstones, and even sandstones. Exogastric shells are thick-walled, with rather variable outer sculpture due to age, location on shell, and preservation. Dorsal, exhalant V-shaped sinus and two inhalant sinuses are present near the suture. Size max. about 35 mm. Many specimens yielded internal moulds with well-preserved retractor muscle scars. The maximum of distribution is in the Záhořany Formation. Specimens from the Libeň and Letná Formations, generally poorly preserved in sandstones, may belong to a separate species. *S. neglecta* probably was a semiinfaunal deposit feeder which lived with the soft body shallowly buried in the surface layer of sediment. One specimen found in the Záhořany Formation shows numerous repaired fractures due to repeated and failed predaceous attacks, probably by nautiloids. Reported from extra-Barrandian territories (Baltic area – Dzik 1981; Morocco – Horný 1997g).

Stratigraphy, localities. Lower part of the Upper Ordovician, Berounian, Libeň to Bohdalec Formations, most common in the Záhořany F.: Beroun, Lodenice, Praha – Jinonice, Praha – Sporilov, Praha – Štěrboholy.

Sinuitopsis hornyi MAREK, 1963

Pl. 1, figs 5, 6

- 1963 *Sinuitopsis hornyi* n. sp.; Marek, Bellerophontid gastropods...
 1963b *Sinuitopsis hornyi* MAREK; Horný, Lower Paleozoic Bellerophontina...

A rare species, found in decalcified siltstones of the uppermost parts of the Kosov Formation. It distinguishes from the type species by weaker, thinner growth structures, wider shell, and narrower umbilicus. Size max. about 20 mm. All types and additional material are deposited in the Czech Geological Survey.

Stratigraphy and locality. Upper Ordovician, Kosovian, Kovsov F.: Praha – Běchovice.

Sinuitopsis sp.

- 1997c *Sinuitopsis* sp.; Horný, New, rare, ...

A fragment of the final whorl, 18 mm long, with patches of external mould with remains of outer sculpture consisting of dense ribs indicating a deep, V-shaped, dorsal sinus. It may belong to *Sinuitopsis evoluta*.

Stratigraphy, locality. Middle Ordovician, Dobrotivian, Dobrotivá F.: Praha – Šárka.

Sinuitopsis HORNÝ, 1997

Type species: *Sinuitopsis plasi* HORNÝ, 1997. Middle Ordovician, Bohemia.

Sinuitopsis plasi HORNÝ, 1997

Pl. 1, figs 7, 8

- 1967c *Sinuitopsis plasi* sp. n.; Horný, new, rare, ...

Species known from a single, incomplete specimen. Nevertheless, characters of the outer shell sculpture clearly show a deep V-shaped sinus at rounded dorsum, and dense, fine, com marginal threads crossed with dense, fine spiral ribs. These features are unknown in other genera. The fragment is 18.5 mm long.

Stratigraphy, locality. Middle Ordovician, Dobrotivian, Dobrotivá F.: Praha – Šárka.

Cyrtodiscus PERNER, 1903

Type species: *Oxydiscus (Cyrtodiscus) procer* BARRANDE in PERNER, 1903. Upper Ordovician, Bohemia.

Cyrtodiscus nitidus

(BARRANDE in PERNER, 1903)

Pl. 1, figs 9, 10

- 1903 *Oxydiscus (Cyrtodiscus) nitidus* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1962 *Gamadiscus nitidus* (PERNER, 1903); Horný, New genera....
 1963b *Gamadiscus nitidus* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
 1997g *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903); Horný, Ordovician Tergomya...

A species with characteristic, sharp, cancellate outer shell sculpture. Spiral ribs continuous. It is common in siliceous concretions, chiefly in the Rokycany region. A cyrtonellid muscle scar pattern was found in one internal mould. Several juveniles yielded fossilized intestinal contents *in situ*. Adult specimens probably lived on the bottom, while the juveniles supposedly inhabited floating algae. Max. size about 10 mm. Reported from the Llanvirnian of Morocco.

Stratigraphy and localities. Middle Ordovician, Llanvirnian, Šárka F.; Osek (common), Praha – Šárka. Dobrotivian, Dobrotivá F.; Malé Přílepy (sporadic).

Cyrtodiscus procer

(BARRANDE in PERNER, 1903)

Pl. 1, figs 11-13

- 1903 *Oxydiscus (Cyrtodiscus) procer* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1941 *Cyrtodiscus procer* (PERNER, 1903); Knight, Paleozoic gastropod genotypes
 1997c *Cyrtodiscus procer* (PERNER, 1903); Horný, New, rare ...

A scarce species, known from two Barrande's types and several fragmentary specimens, collected by L. Marek. Flat,

discoidal shells with narrow periphery bear strong, regular spiral ribs crossed with fine collateral lines. All specimens come from siltstones and silty sandstones. Size up to 20 mm.

Stratigraphy, localities. Upper Ordovician, Berounian, Letná F.: Trubská, Praha – Vysočany.

Cyrtodiscus simaki HORNÝ, 1997

Pl. 1, fig. 14

1997c *Cyrtodiscus simaki* sp. n.; Horný, New, rare, ...

Shell similar to *C. procer* but less regular; weak spiral elements of outer shell sculpture slightly discontinuous in ontogenetically older stages. Size max. up to 10 mm. Rare.

Stratigraphy, localities. Upper Ordovician, Berounian, Záhořany F.: Praha – Spořilov, Praha – Nový Hloubětín, Praha – Dubec.

Cyrtodiscus? evolvens

(BARRANDE in PERNER, 1903)

Pl. 1, fig. 15

1903 *Cyrtolites? evolvens* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1

1963 *Temnodiscus evolvens* (PERNER, 1903); Marek, Bellerophontid gastropods...

1963b *Temnodiscus? evolvens*; Horný, Lower Palaeozoic Bellerophontina...

Shell flat, narrow at the periphery, with regular transverse ribs and discontinuous, weaker spiral ribs. It distinguishes from biostratigraphically older representatives of *Cyrtodiscus* by discontinuous spiral elements in outer shell sculpture. Maximum length 10 mm. Rare, usually flattened in shales. The generic determination is uncertain, as the continuity or discontinuity of spiral ribs seem to be not too reliable systematic feature. *Temnodiscus* KOKEN, 1896 has the final whorl out of contact and its transverse ribs are much coarser.

Stratigraphy and localities. Upper Ordovician, Královorian, Králův Dvůr F.: Králův Dvůr – Kosov, Praha – Radotín. Kosovian, Kosov F.: Praha – Běchovice.

Family *Carcassonnellidae* HORNÝ, 1997

Carcassonnella HORNÝ et PEEL, 1996

Type species: *Gamadiscus courtoissolei* YOCHELSON, 1882. Lower Ordovician, France.

Carcassonnella pragensis HORNÝ, 1997

Pl. 2, figs 1, 2

1997c *Carcassonnella pragensis* sp. n.; Horný, New, rare, ...

Almost globose shells with fine growth lines; selenizone on uniformly rounded dorsum. Size max. 20 mm. Rare, so far found at Prague.

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Praha – Šárka.

Carcassonnella sp.

1997c *Carcassonnella* sp.; Horný, New, rare, ...

Small globose shells, outer sculpture unknown. Size up to 10 mm. Commonly in ferrolites.

Stratigraphy, locality. Middle Ordovician, Dobrotivian, Dobrotivá F.: gallery Veronika near Kařízek.

Sarkanella HORNÝ, 1997

Type species: *Plectronotus? vokovicensis* HORNÝ, 1963. Middle Ordovician, Bohemia.

Sarkanella vokovicensis (HORNÝ, 1963)

Pl. 2, figs 3, 4

1963b *Plectronotus? vokovicensis* n. sp.; Horný, Lower Paleozoic Bellerophontina...

1997c *Sarkanella vokovicensis* (HORNÝ, 1963); Horný, New, rare ...

Small, flat, densely transversely ribbed shells with zigzagging ribs on lateral sides; dorsum with wide selenizone; size up to 14 mm.

Stratigraphy, localities. Middle Ordovician, Dobrotivian, Dobrotivá F.: Mýto, Praha – Vokovice, Praha – Holešovičky (siliceous concretions, not common); Plzenec (clayey shales, common).

Sarkanella novaki HORNÝ et VONKA, 2002

Pl. 2, figs 5, 6

2002 *Sarkanella novaki* sp. n.; Horný and Vonka, *Sinuites* community...

Rare, known from two specimens found in clayey shales, in a taphocoenose with *Sinuites* sp. Similar to *S. vokovicensis* but with wider selenizone. Size about 16 mm.

Stratigraphy, locality. Upper Ordovician, Berounian, Bohdalec Fm.: Praha – Radotín.

Sarkanella kolebabai HORNÝ, 2002

Pl. 2, fig. 7

2002a *Sarkanella kolebabai* sp. n.; Horný, The youngest representative...

The biostratigraphically youngest representative of the genus, known from a unique specimen from clayey shale. It distinguishes by sparse, on the lateral sides weakly zigzagging ribs. Size 9 mm.

Stratigraphy, locality. Upper Ordovician, Královorian, Králův Dvůr F.: Praha – Pankrác.

Peelerophon YOCHELSON, 1982

Type species: *Bellerophon oehlerti* BERGERON, 1889. Lower Ordovician, France.

Peelerophon mergli FRÝDA, 1988

Pl. 2, figs 8, 9

1988 *Modestospira mergli* sp. n.; Frýda, A new species...

1997c *Peelerophon mergli* (FRÝDA, 1988); Horný, New, rare, ...

Small, irregularly ribbed shell, rounded dorsum either with a narrow sinus or selenizone. Size about 11 mm. The species rarely occurs in redeposited, reddish tuffs. The type specimens have been housed in the Czech Geological Survey, Prague.

Stratigraphy, localities. Lower Ordovician, Arenigian, top of the Klabava F.: Ejpovice, Strašice, Kleštěnice.

GASTROPODA

Iosostrophic gastropods do not represent a natural, homogeneous systematic unit. As their systematic is so far not satisfactorily resolved, the suprafamiliar taxa are not given. For the purpose of this catalogue, also patelliform gastropods are included, often classified as "monoplacophorans" or groups of uncertain position. Of these, *Archinacella* ULRICH et SCOFIELD, 1897 and its allies have been already recognized as gastropods (Yochelson 1988, Horný 1996c, Peel and Horný 1999), but the position of many patelliform shells still remain unsolved. While *Floripatella rousseau* YOCHELSON, 1988 possesses a typical patellid muscle scar pattern, other Ordovician (and, of course, also Cambrian) limpet-like shells either lack muscle scars or have them presumably arranged in a more or less continuous circle, parallel or subparallel to the growth structures. This is e.g. the case of *Patelliconus* HORNÝ, 1961, assigned by Geyer (1994) to the Family Protoconchoiidae GEYER, 1994 (together with *Archinacella* and *Floripatella* YOCHELSON, 1988). This muscle scar pattern (as well as the shell shape) is a result of the stationary mode of life on various hard objects and shells on the sea bottom, and may have simultaneously appeared in diverse lineages. An excellent example of such adaptation is the Middle Ordovician *Pygmaeoconus porrectus* (BARRANDE in PERNER, 1903), adjusted to the dorsal surface of a hyolith shell. This species yielded the best-preserved muscle scars in this group of gastropods.

Sinuitids are well-known bellerophontoideans, distributed throughout the whole Ordovician but most abundant in its middle part. They were probably semiinfaunal predators, living in soft, often muddy sediment (Horný 1996a). *Sinuites* KOKEN, 1896 is the first bellerophontid, in which a pair of circumbilical muscles was ascertained (Knight 1947). This pattern, generally accepted as a characteristic sign for endogastric gastropod shells, was later found in additional bellerophontoideans, e.g. *Tremantotus* HALL, 1865, *Megalomphala* ULRICH et SCOFIELD, 1897, *Bellerophon* MONTFORT, 1808, and *Carinaropsis* HALL, 1847 (Peel 1972, 1976, 1982, 1993), including several genera from the Ordovician of Bohemia (*Tropidodiscus* MEEK et WORTHEN, 1866, *Bucanopsis* HORNÝ, 1997, *Tritonophon* ÖPIK, 1953, and *Grandostoma* HORNÝ, 1962).

Bucaniidae are represented by the genus *Bucanopsisina*, quite common in Upper Ordovician, and the rare *Tritonophon*. It is remarkable that the widely distributed genera *Bucania* HALL, 1847, *Salpingostoma* ROEMER, 1876, and *Megalomphala* ULRICH in ULRICH et SCOFIELD, 1897 are absent in Bohemia.

Tropidodiscus is the dominant Middle Ordovician species in Bohemia. *T. pusillus*, which probably lived on floating algal carpets, occurs in innumerable masses; nevertheless, the genus is almost absent in the Upper Ordovician. *Pterothecidae* occur rarely in the Upper Ordovician. Representatives of *Cymbularia*, widely distributed in the Baltic area, are rare, limited to the Middle Ordovician.

Characteristic and common component of the Upper Ordovician benthic fauna are species of *Grandostoma*. *G. grande* is an example of rare immigrants from the Balto-Scandinavian region (Ebbestad 1999).

Of the extra-Barrandian gastropods, types and referred specimens of Horný's Morocco paper (1997g) are deposited in the National Museum, Prague (genera *Sinuites*, *Lesinuites*, *Tritonophon*, *Bucanopsisina*, *Tropidodiscus*, and *Atlantophon* HORNÝ, 1967). Besides these, a single specimen of *Conradella* [= *Bucania*] *gracillima* (KOKEN, 1896) from Dalarne (inv. No S 2439) is also deposited here, which is the type of Koken and Perner 1925, Pl. 27, figs 21-24.

Family *Archinacellidae* KNIGHT, 1956

Barrandicella PEEL et HORNÝ, 1999

Type species: *Archinacella ovata* BARRANDE in PERNER, 1903; Middle Ordovician, Bohemia.

Barrandicella ovata

(BARRANDE in PERNER, 1903)

Pl. 2, figs 10-12

- 1903 *Archinacella ovata* var. *typica* [= *ovata*] BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
1903 *Archinacella ovata* var. *complanata* m.; Perner, Système silurien, IV, Gastéropodes, 1
1903 *Archinacella ovata* var. *rostrata* m. (partim): Perner, Système silurien, IV, Gastéropodes, 1
1903 *Archinacella ovata* var. *subglobosa* m.; Perner, Système silurien, IV, Gastéropodes, 1
1963a *Archinacella ovata* PERNER, 1903; Horný, Lower Paleozoic Monoplacophora...
1996c *Archinacella ovata* BARRANDE in PERNER, 1903; Horný, Apical muscle...
1999 *Barrandicella ovata* (BARRANDE in PERNER, 1903); Peel and Horný, Muscle scars...

A common species, occurring in siliceous concretions of the Middle Ordovician formations. Perner (1903) named several varieties; two of them have persisted (var. *elevata* and var. *tarda*), but var. *complanata* and var. *suglobosa* are inaccurately defined and hardly distinguishable. Related species occur in the Ordovician of the Mediterranean and Baltoscandian regions, England, North America, Greenland, and Asiatic Russia. Specimens reported from the Montagne Noire, assigned to as *A. ovata* (Yochelson 1982), belong to a different, probably not yet described species. In 1996, a pair of retractor muscle scars was ascertained in the apical area of *B. ovata*; this feature clearly demonstrated that *Barrandicella* and its allies were not tergomyans (Horný 1996c, Peel and Horný 1999). *Barrandicella ovata* is the most common Bohemian archinacelloid species, occurring in siliceous concretions in taphocoenoses of other invertebrate fauna. Some concretions contain assorted accumulations of *Barrandicella* shells. It does not occur in sandy shallow water sediments and in deeper water clayey deposits, obviously because of anoxic environment, lack of suitable hard objects, and food in this facies. *B. ovata* was probably a slowly moving grazer feeding on algal mats

overgrowing various shells and fragments lying on the bottom. Size: up to 20 mm. So far, several hundred specimens were collected, which are predominantly deposited in the collections of the Museum of Dr. B. Horák at Rokycany and the National Museum, Prague.

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F.: Osek, Praha – Šárka, Úvaly. Dobrotivian, Dobrotivá F.: Malé Přílepy, Praha – Šárka.

***Barrandicella holubi* HORNÝ, 2000**

Pl. 2, figs 13, 14

2000 *Barrandicella holubi* sp. n.; Horný, *Barrandicella holubi*...

A rare species, similar to *B. ovata* but with narrower shell and a wide, bulky apex. It was found only in the Rokycany region.

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Osek.

***Barrandicella?* *tarda* (PERNER, 1903)**

Pl. 3, figs 3, 4

1903 *Archinacella ovata* var. *tarda* m.; Perner, Système silurien, IV, Gastéropodes, 1

1963a *Archinacella tarda* PERNER, 1903; Horný, Lower Paleozoic Monoplacophora...

1999 *Barrandicella tarda* (PERNER, 1903); Peel and Horný, Muscle scars...

A rare species, sporadically occurring in silty, calcareous concretions. It is relatively most common in the Zahořany Formation in the territory between Beroun and Praha – Štěrboholy, from where we know about ten specimens. It rarely occurs in shales, usually together with remains of other fauna. An adult shell is less globose than in *B. ovata*, the apical part is more protruding, and the concentric lines of growth are crossed with fine, dense radial threads. Presence of this type of sculpture is unusual among archinacelloids (*Helcionopsis* ULRICH et SCOFIELD, 1897), and may indicate a separate genus. Several specimens resembling this species were found in siliceous concretions from the Dobrotivá Formation near the village of Malé Přílepy (collections of C. Klouček and F. Hanuš). They differ from *B. ovata* by less arched shells and less perpendicular posterior shell margin.

Stratigraphy, localities. ?Middle Ordovician, Dobrotivian, Dobrotivá F.: Malé Přílepy. Upper Ordovician, Berounian, Vinice to Bohdalec Fs: Loděnice, Praha – Dubeč, Praha – Štěrboholy.

***Barrandicellosis* HORNÝ, 2000**

Type species: *Archinacella ovata* var. *elevata* PERNER, 1903. Middle Ordovician, Bohemia.

***Barrandicellosis elevata* (PERNER, 1903)**

Pl. 3, figs 1, 2

1903 *Archinacella ovata* var. *elevata*; Perner, Système silurien, IV, Gastéropodes, 1

1903 *Archinacella ovata* var. *rostrata* m. (*partim*); Perner, Système silurien, IV, Gastéropodes, 1

- 1963a *Archinacella? ovata elevata* PERNER, 1903; Horný, Lower Paleozoic Monoplacophora...
- 1963c *Cyrtonellopsis elevata* (PERNER, 1903); Horný, On the systematic...
- 1999b *Barrandicellosis elevata*; Horný, Gastropod...

A rare species, so far found in siliceous concretions in the Rokycany region. It differs from *Barrandicella* by much higher adult shell, projecting apex, and a thick shell wall. Outer shell sculpture consists of simple growth lines, muscle scars are similar to those in *Barrandicella*. Length up to 22 mm. The majority of specimens (about 10) are deposited in the Museum of Dr B. Horák at Rokycany.

Stratigraphy and localities. Middle Ordovician, Llanvirnian, Šárka F.: Osek, Kařez.

Barrandicellosis?* *extenuata

(BARRANDE in PERNER, 1903)

Pl. 2, figs 5, 6

- 1903 *Orthonychia extenuata* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
- 1911 *Orthonychia? extenuata* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 3
- 2000a *Barrandicellosis?* *extenuata* (BARRANDE in PERNER, 1903); Horný, *Barrandicellosis*, ...

The species was based on a single internal mould from sandstones of the Letná Formation at Děd near Beroun. Three additional internal moulds were found quite recently in fields between Trubská and Trubín (I. Chlupáč, Š. Rak). Imperfectly preserved internal moulds do not allow exact generic determination.

Stratigraphy, localities. Upper Ordovician, Berounian, Letná F.: Beroun, Děd; fields between Trubská and Trubín.

***Marekicella* HORNÝ, 1997**

Type species: *Marekicella podexpulicis* HORNÝ, 1967. Upper Ordovician, Bohemia.

***Marekicella podexpulicis* HORNÝ, 1997**

Pl. 3, figs 7, 8

1997c *Marekicella podexpulicis* sp. n.; Horný, New, rare, ...

Rare archinacellid species with a narrow, laterally compressed shell, external shell surface with fine growth lines. Muscular structures are imperfectly known. Max. length 8 mm. The animal may have lived in dense algal growths or on algae.

Stratigraphy, localities. Upper Ordovician, Berounian, Zahořany to Bohdalec F.: Praha – Radotín, Staňkovka; Praha – Vršovice.

***Archinacellina* HORNÝ, 1961**

Type species: *Archinacella modesta* BARRANDE in PERNER, 1903; Upper Ordovician, Bohemia.

Archinacellina modesta

(BARRANDE in PERNER, 1903)

Pl. 3, fig. 9

- 1903 *Archinacella modesta* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1

- 1903 *Archinacella proxima* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1903 *Archinacella capuloidea* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
 1903 *Archinacella mitra* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
 1961 *Archinacellina modesta* (PERNER, 1903); Horný, New genera...
 1963a *Archinacellina modesta* (PERNER, 1903); Horný, Lower Paleozoic...
 1999 *Archinacellina modesta* (BARRANDE in PERNER, 1903); Peel and Horný, Muscle scars...

Relatively rare species, which is characteristic for clayey shales. Most of the collected specimens (about 30) come from the Barrande's collection and are housed in the National Museum, Prague. Outer shell sculpture is unknown, but two internal moulds yielded remains of a loop-like muscular impression, in the posterior part of the shell widened into two subtriangular, equilateral extensions. Outer shell sculpture unknown.

Stratigraphy, localities. Upper Ordovician, Královorianský, Králův Dvůr F.: Chodouň, Beroun – Králův Dvůr.

Family ? *Archinellidae* KNIGHT, 1956
 (? *Protoconchiodidae* GEYER, 1994)

***Patelliconus* HORNÝ, 1961**

Type species: *Palaeacmaea primula* BARRANDE in PERNER, 1903. Middle Ordovician, Bohemia.

Patelliconus primulus
 (BARRANDE IN PERNER, 1903)

Pl. 3, figs 10, 11

- 1903 *Palaeacmaea primula* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1903 *Palaeacmaea immigrans* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1961 *Patelliconus primulus* (PERNER, 1903); Horný, New genera...
 1963a *Patelliconus primulus* (PERNER, 1903); Horný, Lower Paleozoic Monoplacophora...

A small, low, cone-like mollusc with planar aperture, variable in shape, rarely occurring in Middle Ordovician siliceous concretions. Shells, preserved as internal moulds, usually bear a concentric band probably connected with musculature. The animal may have been almost sessile, attached to various shells or fragments, overgrown with algal or bacterial film.

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F.: Osek, Praha – Šárka. Dobrotivian, Dobrotivá F.: Mýto, Praha – Šárka.

***Pygmaeoconus* HORNÝ, 1961**

Type species: *Palaeacmaea porrecta* BARRANDE in PERNER, 1903. Middle Ordovician, Bohemia.

Pygmaeoconus porrectus
 (BARRANDE in PERNER, 1903)

Pl. 4, fig. 1

- 1903 *Palaeacmaea? porrecta* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1961 *Pygmaeoconus porrectus* (PERNER, 1903); Horný, New genera...
 1963a *Pygmaeoconus porrectus* (PERNER, 1903); Horný, Lower Paleozoic...

A small cone-like shell with subcentral apex, common in siliceous concretions of the Šárka Formation but sporadically occurring also in the Dobrotivá Formation. It is usually overlooked because of its small size (max. length 5.8 mm). Internal moulds often show muscle scars arranged in a more or less continuous band encircling the shell. This pattern is connected with the presumed stationary habit of *Pygmaeoconus*. Moreover, the aperture was not planar but concavely bent and thus adapted to upper, convex, dorsal surface of a hyolithid shell, which these tiny molluscs probably inhabited. *Pygmaeoconus* either grazed organic film on the host shell or filtrated deposit, swirled up by the hyolith when feeding (Horný, a paper in preparation).

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F.: Osek, Praha – Šárka. Dobrotivian, Dobrotivá F.: Mýto, Praha – Šárka.

***Pygmaeoconus kettneri* (ŘÍHA, 1938)**

Pl. 3, fig. 12

- 1938 *Palaeacmaea kettneri* n. sp.; Říha, Příspěvek k poznání...
 1963a *Pygmaeoconus? kettneri* (ŘÍHA, 1938); Horný, Lower Paleozoic Monoplacophora...

A rare species with small (up to 7.5 mm), cap-shaped shell with subcentral apex; external surface with simple growth lines. May be conspecific with *P. porrectus*.

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Osek.

Pygmaeoconus? latiusculus
 (BARRANDE in PERNER, 1903)

Pl. 4, fig. 7

- 1903 *Palaeacmaea latiuscula* Barr. sp.; Perner, Système silurien, IV, Gastéropodes, 1
 1963a *nomen dubium*; Horný, Lower Paleozoic Monoplacophora...

Imperfectly preserved internal mould, 5.6 mm long, does not allow satisfactory description and determination.

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Osek.

***Pygmaeoconus krafti* HORNÝ, 1997**

Pl. 4, figs 2, 3

- 1997c *Pygmaeoconus krafti* sp. n.; Horný, New, rare, ...

Rare, small (up to 8.5 mm), elongate cap-shaped shell with subcentral apex and irregular muscle scars located in a ring-shaped area parallel with the apertural margin. Outer shell sculpture with irregularly spaced, fine, concentric lines of

growth. Probably less mobile, living on hard objects like *P. porrectus*.

Stratigraphy, localities. Upper Ordovician, Berounian, Zahroňany F.; Lodenice, Praha – Libeň, Praha – Hloubětín, Praha – Dubeč. Bohdalec F.: Praha – Michle.

? *Micropileus* WILSON, 1951

Type species: *Micropileus obesus*, WILSON, 1951. Middle Ordovician, Canada.

Micropileus? *ordovicinus* (HORNÝ, 1963)

Pl. 4, fig. 4

1963a *Palaeoscurria ordovicina* sp. n.; Horný, Lower Paleozoic Monoplacophora...

A cap-shaped shell, in gross morphology similar to the Silurian *Palaeoscurria calyptrata* PERNER, 1903, and Middle Ordovician *Micropileus variabilis* WAHLMAN, 1992. The only existing specimen, an incomplete, weathered internal mould, is rather insufficiently preserved, and the subapical muscle scar, mentioned by Horný, should be re-evaluated. Size 10 mm. Until better preserved specimens are found, I prefer to assign this species, although with a question, to the Ordovician genus *Micropileus* WILSON, 1951.

Stratigraphy, locality. Middle Ordovician, Dobrotivian, Dobrotivá F.: Malé Přílepy.

Kornoutella HORNÝ, 1997

Type species: *Hypseloconus?* *bohemicus* HORNÝ, 1963. Middle Ordovician, Bohemia.

Kornoutella bohemica (HORNÝ, 1963)

Pl. 4, figs 5, 6

1963a *Hypseloconus?* *bohemicus* sp. n.; Horný, Lower Paleozoic Monoplacophora...

1997c *Kornoutella bohemica* (HORNÝ, 1963); New, rare, ...

Two specimens of this relatively large, cornet-like, about 12 mm high, radially ribbed fossil with thick, probably two-layered shell were so far found in siliceous concretions. Its detailed morphology is unknown and a peculiar outer shell sculpture is unusual among other similar molluscs. Its systematic position is uncertain.

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Praha – Šárka.

Family *Sinuitidae* DALL in ZITTEL-EASTMAN, 1913

Sinuites KOKEN, 1896

Type species: *Bellerophon bilobatus* SOWERBY, 1839. Middle Ordovician, England.

Sinuites sp.

The collection of sinuitids deposited in the Museum of Dr B. Horák at Rokycany includes several flattened and deformed moulds of *Sinuites* sp., coming from shales of the Lower

Ordovician Klabava Formation. Specimens mentioned by Holub (1911, 1912) are also in this collection. Details of external morphology are lacking, but the species is not conspecific with *S. sowerbyi* PERNER, 1903.

Stratigraphy, locality. Lower Ordovician, Arenigian, Klabava F.: Klabava.

Sinuites sowerbyi PERNER, 1903

Pl. 4, figs 8-10

- 1903 *Sinuites Sowerbyi* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
1903 *Sinuites Sowerbyi* var. *crenata* mihi; Perner, Système silurien, IV, Gastéropodes, 1
1963b *Sinuites sowerbyi* PERNER, 1903; Horný, Lower Paleozoic Bellerophontina...
1992 *Sinuites sowerbyi* PERNER, 1903; Horný, Svalové vltisky...
1996a *Sinuites sowerbyi* PERNER, 1903; Horný, Secondary shell deposits...
non 1903 *Sinuites Sowerbyi* var. *evoluta* mihi; Perner, Système silurien, IV, Gastéropodes, 1

The most common sinuitid gastropod of the Bohemian Middle Ordovician, known from many hundreds of specimens preserved in siliceous concretions and clayey shales. The biggest specimens reach about 30 mm. External sculpture consists of relatively coarse transversal ribs; a characteristic dorsal, symmetrically branching sculpture of unclear origin is an important determination feature. Numerous internal moulds yielded well-preserved retractor muscle scars. Two specimens were found showing small patches of inductural deposits. The species occurs also in the Llanvirnian strata in Morocco (Horný 1997g) but specimens from the Arenigian formations of the Montagne Noire, reported by Yochelson (1982) as *Sinuites* cf. *sowerbyi*, represent a different species. Internal moulds cannot be reliably distinguished from those of *Sinuites hanusi* and *Selesinuites perneri*. *S. sowerbyi* has been reported from the Middle Ordovician of Morocco (Horný 1997g).

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F., common: Osek, Praha – Šárka, Praha – Libuš, Úvaly; Dobrotivian, Dobrotivá Formation, rarer: Malé Přílepy, Praha – Šárka.

Sinuites hanusi HORNÝ, 1997

Pl. 5, fig. 6

1997c *Sinuites hanusi* sp. n.; Horný, New, rare, ...

A species similar to *Sinuites sowerbyi* (PERNER, 1907). It distinguishes by finer and denser collateral ribs and slightly narrower shell. It is less frequent than *S. sowerbyi*. It occurs in siliceous concretions, rather in Prague and Úvaly than in the Rokycany region. Internal moulds are hardly distinguishable from those of *S. sowerbyi*. An internal mould of *Sinuites* sp. from Osek (MBHR 1232) with a well-preserved muscle scar, figured by Horný (1992, Pl. 6 and Text-fig. 7), may belong to this species.

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F.: Osek, Praha – Šárka, Úvaly.

Sinuites reticulatus PERNER, 1903

Pl. 4, fig. 11; Pl. 5, figs 1-3

- 1903 *Sinuites reticulatus* PERNER; Perner, Système silurien, IV, Gastéropodes, 1

- 1963b *Sinuites reticulatus* PERNER, Horný, Lower Paleozoic Bellerophontina...
 1992 *Sinuites reticulatus* PERNER, 1903; Horný, Svalové vtisky...
 1996a *Sinuites reticulatus* PERNER, 1903; Horný, Secondary shell deposits...

A common Bohemian Middle Ordovician sinuitid gastropod, known from several hundred specimens preserved in siliceous concretions. The largest specimens reach about 40 mm length. External sculpture consists of dense, fine, sharp spiral and collabral lirae. Numerous internal moulds yielded well-preserved retractor muscle scars. About 30 specimens from Cheznovice yielded exclusively preserved inductural deposits and inductural vallum fossilized in SiO_2 (Horný 1996a), which contributed to explain the mode of life of sinuitids.

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F. (common): Osek, Kařízek – Pětidomky, Praha – Šárka, Úvaly; Dobrotivian, Dobrotivá F. (rare): Malé Přílepy, Praha – Šárka.

Sinuites sp.

- 2002 *Sinuites* sp.; Horný and Vonka, *Sinuites* community...

Several tens of crushed and flattened internal and composite moulds were found in dark grey clayey shales of the Bohdalec Formation. Preservation is too poor to restore the original sculpture, consisting of dense collabral ribs, probably crossed with dense, fine, spiral threads. Maximum size (estim.) is 33 mm. The find is important, demonstrating a continuous presence of *Sinuites* communities since Arenigian till Královorian in the Ordovician of Bohemia.

Stratigraphy, locality. Upper Ordovician, Berounian, Bohdalec F.: Praha – Radotín.

Sinuites aff. *bilobatus* (SOWERBY, 1839)

Pl. 5, fig. 5

- 1903 *Sinuites bilobatus* SOWERBY var. *infaustus* BARR. sp.; Perner, Système silurien, Gastéropodes, IV, 1
 1963b *Sinuites* aff. *bilobatus* (SOWERBY, 1839); Horný, Lower Paleozoic Bellerophontina...

A rare sinuitid known from two poorly preserved, incomplete internal moulds, and several flattened specimens. The taxon cannot be exactly described and its satisfactorily assignment to the British species is doubtful. External shell sculpture consists of simple, fine, collabral ribs.

Stratigraphy, locality. Upper Ordovician, Královorian, Králov Dvůr F.: Chodouň, Beroun – Králov Dvůr.

Sinuites strangulatus (BARRANDE in PERNER, 1903)

Pl. 5, fig. 4

- 1903 *Bellerophon* (*Sinuites?*) *strangulatus* BARR.; Perner, Système silurien, IV, Gastéropodes, 1
 1962 *Strangulites strangulatus* (PERNER, 1903); Horný, New genera...
 1963b *Strangulites strangulatus* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
 1990 *Sinuites* (*Strangulites*) *strangulatus* (PERNER, 1903); Horný, Svalové vtisky...
 1991 *Sinuites* (*Strangulites*) *strangulatus* (PERNER, 1903); Horný, Další poznatky...

A rare sinuitid from the Králov Dvůr Formation with large (up to 30 mm), almost spherical shells, usually flattened in shales. Outer shell sculpture finely cancellate. Exceptionally well-preserved internal moulds from carbonate concretions at the Levín Hill yielded retractor muscle scars. Classification as a separate genus or a subgenus of *Sinuites* has not been affirmed (e.g. Wahlman 1992).

Stratigraphy, localities. Upper Ordovician, Královorian, Králov Dvůr F.: Chodouň, Beroun – Králov Dvůr, Levín.

Selesinuites HORNÝ, 1997

Type species: *Selesinuites perneri* HORNÝ, 1967. Middle Ordovician, Bohemia.

Selesinuites perneri HORNÝ, 1967

Pl. 5, figs 7, 8

- 1967c *Selesinuites perneri* sp. n.; Horný, New, rare, ...

A rare sinuitid similar to *S. sowerbyi* but with a low, wide keel with a pseudoselenizone in immature shells. Outer shell sculpture, dense collabral ribs; remains of subinductural deposits observed in the holotype. This species was reported from the Middle Ordovician of Morocco (Horný 1997g).

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Osek.

Family *Tropidodiscidae* KNIGHT, 1956

Tropidodiscus MEEK et WORTHEN, 1866

Type species: *Bellerophon curvilineatus* CONRAD, 1842. Lower Devonian, USA.

Tropidodiscus pusillus

(BARRANDE in PERNER, 1903)

Pl. 1, fig. 10; Pl. 5, fig. 9

- 1903 *Temnodiscus pusillus* BARR., Perner, Système silurien, IV, Gastéropodes, 1
 1962 *Tropidodiscus* (*Peruniscus*) *pusillus* (PERNER, 1903); Horný, New genera...
 1963b *Tropidodiscus* (*Peruniscus*) *pusillus* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
 1997c *Tropidodiscus pusillus* (BARRANDE in PERNER, 1933); Horný, New, rare, ...
 1997g *Tropidodiscus pusillus* (BARRANDE in PERNER, 1933); Horný, Ordovician Tergomya...

The most common bellerophontidean gastropod in the Ordovician of Bohemia, which occurs in masses in Middle Ordovician siliceous concretions and clayey shales. The sculpture consists of fine growth lines. Its shells are usually small (in average about 3–5 mm), but rarely occur large specimens reaching nearly 15 mm. The shell wall was usually damaged during early fossilization processes. Peel (1978) expressed an opinion that this species lived on algae, and Horný (1999a) speculated about pseudoplanktic life in floating algal carpets. A few specimens yielded retractor muscle scars. The species is distributed

in Llanvirnian of Morocco (Horný 1997g) and France (Pillet 1992); Pillet assumed (probably rightfully) that this species was conspecific with the British *Tropidodiscus acutus* (SOWERBY, 1839), and reported it from Anjou. The validity of the subgenus *Tropidodiscus* (*Peruniscus*) HORNÝ, 1962 was not affirmed. A new, related species occurs in the Dobrotivá Formation.

Stratigraphy, localities. Middle Ordovician, Llanvirnian, Šárka F. (abundant): Osek, Těžkov, Kařez, Praha – Šárka, Praha – Libuš, Úvaly; Dobrotivian, Dobrotivá F. (less abundant): Plzenec, Mýto, Malé Přílepy, Praha – Holešovičky, Praha – Šárka.

Tropidodiscus bouceki HORNÝ, 1997

Pl. 5, fig. 10

1997c *Tropidodiscus bouceki* sp. n.; Horný, New, rare, ...

To date, a single siliceous concretion was found containing several specimens of this species. It differs from *T. pusillus* by coarsely ribbed, in average much bigger, up to 17 mm long shell.

Stratigraphy, locality. Middle Ordovician, Llanvirnian, Šárka F.: Praha – Šárka.

Tropidodiscus sp.

1997c *Tropidodiscus* sp. A; Horný, New, rare, ...

Several imperfectly preserved specimens, similar to *T. pusillus*, were found in weathered silty carbonate concretions. Size is about 3-5 mm.

Stratigraphy, localities. Upper Ordovician, Berounian, Záhořany F.: Praha – Spořilov, Praha – Dubec.

Tropidodiscus sp.

1997c *Tropidodiscus* sp. B; Horný, New, rare, ...

The only specimen comes from Barrande's collection. An internal mould is preserved in a decalcified concretion. The shell is laterally compressed, with ribbed external surface. Size 3.5 mm.

Stratigraphy, locality. Upper Ordovician, Královorianský, Králov Dvůr F.: Beroun - Králov Dvůr.

Family BUCANOIDAE ULRICH et SCOFIELD, 1897

Bucanopsis HORNÝ, 1997

Type species: *Bucanopsis calypso* PERNER, 1903; Upper Ordovician, Bohemia.

Bucanopsis roemeri (BARRANDE in PERNER, 1903)

Pl. 6, fig. 1

1903 *Carinariopsis Roemeri* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
1963b *Bucanopsis roemeri* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...

Imperfectly known species, occurring in sandstones and siltstones of the Letná Formation. About 25 mm long, low shell has a widely explanate aperture. Outer sculpture consists of distinct spiral and collateral ribs. The animal lived on shallow sandy and silty bottom. The majority of specimens are preserved as deformed, rough internal moulds.

Stratigraphy, localities. Upper Ordovician, Berounian, Letná F.: Trubská, Praha – Vysočany.

Bucanopsis calypso PERNER, 1903

Pl. 6, figs 2-4

- 1903 *Bucanopsis Calypso* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
- ? 1903 *Bucania comata* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1
- 1963b *Bucanopsis calypso* PERNER, 1903; Horný, Lower Paleozoic Bellerophontina...
- ?1963b *Bucanopsis cf. comata* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
- 1997a *Bucanopsis calypso* (PERNER, 1903); Horný, *Bucanopsis* gen. n., ...

A bellerophontoid with widely explanate, up to 20 mm long shell, bearing fine spiral and collateral threads. It distinguishes from a similar *Grandostoma bohemicum* mainly by the presence of a dorsal selenizone and a strong parietal callus near the aperture. The shells often show repaired fractures after attempted predation. Several internal moulds yielded a pair of circumfulcal retractor muscle scars. The animal probably lived in higher energy conditions on shallow, silty bottom. Poorly preserved *Bucanopsis comata* and *B. cf. comata* probably belong to this species. *B. calypso* has been reported from the Upper Ordovician of Morocco (Horný 1997g).

Stratigraphy, localities. Upper Ordovician, Berounian, Vinnice to Bohdalec Fs: Loděnice, Praha – Libeň, Praha – Štěrboholy.

Bucanopsis? sp.

1963b *Bucanopsis* sp.; Horný, Lower Paleozoic Bellerophontina...

Several specimens flattened in clayey shale from the Králov Dvůr Formation, 15-20 mm big, are deposited in the collections of the National Museum, Prague. They were originally labelled as *Bellerophon roemeri*. Preservation does not allow an accurate determination.

Stratigraphy, localities. Upper Ordovician, Královorianský, Králov Dvůr F.: Libomyšl, Beroun – Králov Dvůr, Kosov.

Tritonophon ÖPIK, 1953

Type species: *Kokenospira (Tritonophon) trimetra* ÖPIK, 1953. Lower Silurian, Victoria.

Tritonophon peeli HORNÝ, 1997

Pl. 6, fig. 5

- 1903 *Bucaniella bohemica* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
- 1963b *Bucanella* sp.; Horný, Lower Paleozoic Bellerophontina...
- 1997c *Bucanella?* sp.; Horný, New, rare, ...
- 1997f *Tritonophon peeli* sp. n.; Horný, Circumfulcal retractor...

1997g *Tritonophon peeli* sp. n.; Horný, Ordovician Tergomya...

A rare species, known from clayey and silty deposits. A small, about 5 mm long, trilobed shell bears fine growth threads and a selenizone on a narrow dorsum. One specimen, already known to Barrande, comes from sandy facies at Dražov (Děd). Another internal mould yielded a circumbilical retractor muscle scar. The animal probably lived on algae.

Stratigraphy, localities. Upper Ordovician, Berounian, Letná to Bohdalec Fs: Beroun, Děd; Loděnice, Praha – Řeporyje, Praha – Vysočany, Praha – Hloubětín.

Tritonophon? bohemicus (PERNER, 1903)

Pl. 6, figs 6, 7

- 1903 *Bucaniella bohemica* PERNER; Perner, Système silurien, IV, Gastéropodes, 1
1963 *Bucanella bohemica* (PERNER, 1903); Marek, Bellerophontid Gastropods...
1963b *Bucanella bohemica* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
1974 *Tritonophon? bohemica* (PERNER, 1903); Peel, Systematics, ontogeny...

A rare species; trilobed shell with fine spiral and collabral lines; narrow dorsal sinus (or selenizone?). Size less than 10 mm. Found in siltstone concretions and sandstone. The animal probably lived on algae.

Stratigraphy, localities. Upper Ordovician, Královorians, Králův Dvůr F.: Chodouň, Praha – Běchovice.

Eobucania KOBAYASHI, 1955

Type species: *Eobucania pulchra* KOBAYASHI, 1955. Lower Ordovician, British Columbia.

Eobucania? bohemica HORNÝ, 1997

Pl. 6, fig. 8

1997c *Eobucania? bohemica* sp. n.; Horný, New, rare, ...

A unique, fragmentary, globose shell, about 8 mm big, shows fine collabral threads and a wide selenizone on a rounded dorsum. Assignment of this species to *Eobucania* is, however, tentative and needs study of additional material. The shell shape and a steep angle of growth lines with a surprisingly wide selenizone may indicate a relationship with specialized Lower Ordovician representatives of coiled tergomyans *Thoralispira* HORNÝ et VIZCAÍNO, 1995 and *Carcassonnella* HORNÝ et PEEL, 1996.

Stratigraphy, locality. Middle Ordovician, Dobrotivian, Dobrotivá F.: Praha – Šárka.

Family *Pterothecidae* WENZ, 1938

Pterotheca SALTER, 1853

Type species: *Atrypa transversa* PORTLOCK, 1843; Middle Ordovician, Ireland.

Pterotheca consobrina BARRANDE, 1872

Pl. 6, fig. 9

- 1872 *Pterotheca consobrina* BARR.; Barrande, Système silurien, I, Trilobites, Supplement
-1963b *Pterotheca consobrina* BARRANDE, 1872; Horný, Lower Paleozoic Bellerophontina...

Flat bellerophontoids with reduced spire, narrow dorsal keel and widely expanded apertural margin. The shell has an appearance of a flat brachiopod. Maximum size about 25 mm. *P. consobrina* was adapted to quiet, soft, muddy bottom. Rare; almost all specimens come from Barrande's collection.

Stratigraphy, locality. Upper Ordovician, Královorians, Králův Dvůr F.: Beroun – Králův Dvůr.

Family *Bellerophontidae* MCCOY, 1851

Cymbularia KOKEN, 1896

Type species: *Cymbularia galeata* KOKEN, 1896. Middle Ordovician, Estonia.

Cymbularia klouceki HORNÝ, 1997

Pl. 6, fig. 10

1997c *Cymbularia klouceki* sp. n.; Horný, New, rare, ...

Small, almost globose shells up to 11 mm long, with fine collabral threads and well-developed, low dorsal keel on a rounded dorsum. A characteristic but rare species for the Dobrotivá Formation, occurring in siliceous concretions, rarely in shales. It may have lived pseudoplanktonically, similar to *Tropidodiscus pusillus*.

Stratigraphy, localities. Middle Ordovician, Dobrotivian, Dobrotivá F.: Mýto, Malé Přílepy, Praha – Šárka.

Cymbularia sp.

1997c *Cymbularia* sp.; Horný, New, rare, ...

A unique find from a redeposited concretion in gravel terrace. A keeled shell about 11 mm long, with rhombic transversal section. Stratigraphic position uncertain.

Stratigraphy, locality. Middle Ordovician, Llanvirnian or Dobrotivian, Šárka or Dobrotivá Fs: Praha – Hlubočepy.

Family ? (Subfamily *Grandostomatinae* HORNÝ, 1962)

Grandostoma HORNÝ, 1962

Type species: *Salpingostoma grande* BARRANDE in PERNER, 1903. Upper Ordovician, Bohemia.

Grandostoma bohemicum (PERNER, 1903)

Pl. 6, figs 11, 12

- 1903 *Carinariopsis bohemica* PERNER; Perner, Système silurien, IV, Gastéropodes, 1

- 1903 *Carinariopsis catenularia* PERNER; Perner, Système silurien, IV, Gastéropodes, I
 1903 *Carinariopsis Sardesoni* PERNER; Perner, Système silurien, IV, Gastéropodes, I
 1903 *Bucania delicatula* PERNER; Perner, Système silurien, IV, Gastéropodes, I
 1963b *Grandostoma bohemicum* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
 1996d *Grandostoma bohemicum* (PERNER, 1903); Horný, *Grandostoma*...
 1997b *Grandostoma bohemicum* (PERNER, 1903); Horný, Shell breakage...


An explanate bellerophontacean gastropod, up to 30 mm big, occurring in clayey siltstones and shales of the Berounian stage. It is similar to *Bucanopsis calypso* but lacks selenizone and callus. It was probably adapted to less dynamic conditions but its fossils are often found in taphocoenoses together with *B. calypso*. The best-preserved specimens come from carbonate concretions in basal parts of the Bohdalec Formation in Praha – Hloubětín. Like in *B. calypso*, the shells often show repaired fractures after attempted predation, and some of the internal moulds yielded circumbilical refractor muscle insertions. The species may represent a separate genus.

Stratigraphy, main localities. Upper Ordovician, Berounian, Vincice, Zahořany, and Bohdalec Formations: Loděnice, Praha – Velká Chuchle, Praha – Jinonice, Praha – Libeň, Praha – Štěrboholy.

Grandostoma grande (BARRANDE in PERNER, 1903)

Pl. 6, fig. 13

- 1903 *Salpingostoma grande* BARR. sp.; Perner, Système silurien, IV, Gastéropodes, 1



Ordovician in the central part of the Barrandian Area between Praha and Plzeň (after Chlupáč, I. et al. 1998)

- 1962 *Grandostoma grande* (PERNER, 1903); Horný, New genera...
 1963b *Grandostoma grande* (PERNER, 1903); Horný, Lower Paleozoic Bellerophontina...
 1997e *Grandostoma grande* (BARRANDE in PERNER, 1903); Horný, Circumbilical...

A characteristic species with strong transversal ribs crossed with undulated spiral lines. Spire small. Dorsum with a shallow sinus. One pair of circumbilical retractor muscle insertions. Widely explanate aperture up to 65 mm across testifies adaptation to quiet, soft, clayey bottom. The majority of specimens come from Barrande's collection. According to an opinion of Horný (1963b), confirmed by O. J. Ebbestad (2000), this species is distributed also in the Upper Ordovician of Great Britain.

Stratigraphy, localities. Upper Ordovician, Kralodvorian, Králův Dvůr F: Beroun – Králův Dvůr, Chodouň, Velká Chuchle, Levín.

Grandostoma taconicum MAREK, 1963

Pl. 6, fig. 14

- 1963 *Grandostoma taconicum* n. sp.; Marek, Bellerophontid Gastropods...
 1963b *Grandostoma taconicum* MAREK, 1903; Horný, Lower Paleozoic Bellerophontida...

A rare species known from several incomplete and poorly preserved specimens; size about 15 mm.

Stratigraphy, locality. Upper Ordovician, Kosovian, Kosov Formation; Běchovice near Praha.

Series		lithology		graptolite zones		Tergomya and Gastropoda	
	Formations					Stages	
U P P E R	Berounian	Kralodvorian	Kosovian	Glyptograptus bohemicus		Tergomya: <i>Sinuitopsis hornyi</i> , <i>Cyrtodiscus? evolvens</i> , <i>Sarkanella kolebabai</i> Gastropoda: <i>Trinophon? bohemicus</i> , <i>Grandostoma taconicum</i>	
		Králův Dvůr	Kosov	grey-green claystones	Glyptograptus cf. ojsuensis Scalariograptus angustus Glyptograptus teres	Tergomya: <i>Cyrtodiscus? evolvens</i> Gastropoda: <i>Archinacellina modesta</i> , <i>Sinuites strangulatus</i> , <i>S. aff. bilobatus</i> , <i>Tropidodiscus</i> sp., <i>Bucanopsina?</i> sp. <i>Tritonophon? bohemicus</i> , <i>Pterotheca consobrina</i> , <i>Grandostoma grande</i>	
		Bohdalec	Záhořany	black claystones carbonate silt-stones	Diplograptus vulgatus Dicranograptus cf. clingani	Tergomya: <i>Sinuitopsis neglecta</i> , <i>Sarkanella novaki</i> Gastropoda: <i>Barrandicella?</i> tarda, <i>Marekicella podexpulicis</i> , <i>Pygmaeoconus krafti</i> , <i>Sinuites</i> sp., <i>Bucanopsina calypso</i> , <i>Tritonophon peeli</i> , <i>Grandostoma bohemicum</i>	
		Vinice		siltstones	Glyptograptus insculptus, Glyptograptus trubinensis extensus	Tergomya: <i>Sinuitopsis neglecta</i> , <i>Cyrtodiscus simaki</i> Gastropoda: <i>Barrandicella?</i> tarda, <i>Marekicella podexpulicis</i> , <i>Pygmaeoconus krafti</i> , <i>Tropidodiscus</i> sp., <i>Bucanopsina calypso</i> , <i>Tritonophon peeli</i> , <i>Grandostoma bohemicum</i>	
		Letná		black shales	Glyptograptus trubinensis trubinensis	Tergomya: <i>Sinuitopsis neglecta</i> Gastropoda: <i>Bucanopsina calypso</i> , <i>Grandostoma bohemicum</i>	
		Libeň		sandstones, greywackes, shales		Tergomya: <i>Sinuitopsis neglecta</i> , <i>Cyrtodiscus procer</i> Gastropoda: <i>Barrandicelopsis?</i> extenuata, <i>Bucanopsina roemerii</i> , <i>Tritonophon peeli</i>	
				black shales		Tergomya: <i>Sinuitopsis</i> sp.	
				sandstones		Tergomya: <i>Pentalina prantli</i> , <i>Sinuitopsis</i> sp., <i>Sinuitopsis plasi</i> , <i>Cyrtodiscus nitidus</i> , <i>Carcassonnella</i> sp., <i>Sarkanella vokovicensis</i> Gastropoda: <i>Barrandicella ovata</i> , <i>Barrandicella?</i> cf. <i>tarda</i> , <i>Patelliconus primulus</i> , <i>Micropileus?</i> <i>ordovicinus</i> , <i>Sinuites sowerbyi</i> , <i>S. reticulatus</i> , <i>Tropidodiscus pusillus</i> , <i>?Eobucania bohemica</i> , <i>Cymbularia klouckei</i> , <i>Cymbularia</i> sp	
				black shales	Glyptograptus tricornis	Tergomya: <i>Sinuitopsis evoluta</i> , <i>Cyrtodiscus nitidus</i> , <i>Carcassonnella pragensis</i> Gastropoda: <i>Barrandicella ovata</i> , <i>Barrandicella holubi</i> , <i>Barrandicelopsis elevata</i> , <i>Patelliconus primulus</i> , <i>Pygmaeoconus porrectus</i> , <i>P. kettneri</i> , <i>P.? latiusculus</i> , <i>Kornoutella bohemica</i> , <i>Sinuites sowerbyi</i> , <i>S. reticulatus</i> , <i>S. hanusi</i> , <i>Selesinuites perneri</i> , <i>Tropidodiscus pusillus</i> , <i>T. bouceki</i>	
				greenish shales red deposits	Didymograptus clavulus Corymbograptus retroflexus	Tergomya: <i>Peelerophon mergli</i> Gastropoda: <i>Sinuites</i> sp.	
L O W E R	Tremadocian	Arenigian	Lišovian	Dobrotivian	Tetragraptus abbreviatus Holograptus tardibrachiatus Corymbograptus V-similis (Clonograptus)	Tergomya: <i>Peelerophon mergli</i> Gastropoda: <i>Sinuites</i> sp.	
	Třenice	Mílna	Klabava	Šárka	silicites, red shales		
					sandstones, conglomerates	(Rhabdinopora intermedia)	

Distribution of Tergomya and isostrophic Gastropoda in the Ordovician of Bohemia (partly after V. Havlíček in I. Chlupáč et al. 1998)

CATALOGUE

How to use the catalogue

The catalogue contains all type and referred specimens of Czech Ordovician tergomyans and isostrophic and symmetrical gastropods deposited in the collections of the Department of Palaeontology, Museum of Natural History, National Museum, Prague. The brachiopod *Ptychopeltis incola* PERNER, 1903, originally described as a gastropod (Perner 1903, Pl. 43, figs 8-15), has not been included.

The first line. The items are arranged alphabetically according to specific names, as originally used. For the purpose of the catalogue, all specific names are given with a small initial letter. The specific name precedes the generic, which is separated by a comma. Both names are printed in bold types. In cases that a species was not defined, the generic name is given as first. After the colon, the author/s of taxon with the relevant year is/are given.

The second line gives the year of publication as quoted in the references, its author/s, and an indication of illustration/s. In case that the specimen was figured on a plate and on a text-figure, the

atavum, Salpingostoma?: PERNER, 1903

1903 Perner: Text-fig. 72a, p. 103

OML: Šárka F.; Praha – Lhotka

■L 23777, lectotype (SD Horný herein)

= anorganic structure

atavum, Salpingostoma?: PERNER, 1903

1903 Perner: Text-fig. 72b, p. 103

OML: Šárka F.; Praha – Lhotka

L 23778, paralectotype

= anorganic structure

bicarinatus, Temnodiscus: PERNER, 1903

1903 Perner: Text-figs 52a-d, p. 77

OUN: Letná F.; Trubská

■L 5643, lectotype (SD Horný herein)

T = *Sinuitopsis neglecta* PERNER, 1903 (Horný 1963b)

bicarinatus, Temnodiscus: PERNER, 1903

1903 Perner: AD Text-figs 52a-d, p. 77

OUN: Letná F.; Trubská

L 5644, paralectotype

T = *Sinuitopsis neglecta* PERNER, 1903 (Horný 1963b)

bilobatus var. *infaustus*, *Sinuites*: BARRANDE in PERNER, 1903

1907 Perner: Pl. 112, figs VI/1, 2

OUK: Králov Dvůr F.; Lejškov (= Chodouň)

L 5609, paralectotype

G = *Sinuites* aff. *bilobatus* (SOWERBY, 1839)

bilobatus var. *infaustus*, *Sinuites*: BARRANDE in PERNER, 1903

1907 Perner: Pl. 112, figs VI/3-5; Text-figs 27a-e, p. 60

1963b Horný: Pl. 5, figs 1-3 (*Sinuites* aff. *bilobatus*)

OUK: Králov Dvůr F.; Lejškov (= Chodouň)

■L 5608, lectotype (SD Horný 1963b)

G = *Sinuites* aff. *bilobatus* (SOWERBY, 1839)

bohemica, Bucaniella: PERNER, 1903

1963 Marek: Pl. 2, figs 1-3

1963b Horný: Pl. 9, fig 7 (*Bucanella bohemica*)

1974 Peel: Text-fig. 10, p. 249 (*Tritonophon?* *bohemica*)

OUK: Kosov F.; Praha – Běchovice

L 36542

G = *Tritonophon?* *bohemicus* (PERNER, 1903) (Peel 1974)

plate precedes the figure. In case that the specimen was re-figured, additional line/s is/are intercalated indicating the year, author, illustration/s, and name of the taxon as used. The abbreviation AD means an additional specimen used for the relevant figure.

The third line gives data about the age and locality. After the abbreviation Oxx [OLA (Lower Ordovician, Arenigian), OML (Middle Ordovician, Llanvirnian), OMD (Middle Ordovician, Dobrotivian), OUB (Upper Ordovician, Berounian), OUK (Upper Ordovician, Kralodvorian), and OUko (Upper Ordovician, Kosovian)] follows a name of the formation, and after a semi-colon a name of the locality, both in present usage.

The fourth line contains the catalogue number (L xxxx). In case that the specimen has a counterpart, the symbol ± is added. After comma, the category of nomenclatory type is added; in case of the lectotype, the author and a date of the subsequent designation (SD) are given. The symbol ■ at the beginning of this line indicates the mandatory type (holotype, lectotype).

The fifth line gives an abbreviation of the class name (T – Tergomya, G – Gastropoda), followed by the name in full wording accepted as valid.

bohemica, Bucaniella: PERNER, 1903

1903 Perner: Pl. 86, figs 40-42

1963b Horný: specimen mentioned on p. 81 (*Bucanella* sp.)

OUN: Letná F.; Drabov (= Beroun, Děd)

L 5623, paralectotype

G = *Tritonophon peeli* HORNÝ, 1997 (Horný 1997f)

bohemica, Bucaniella: PERNER, 1903

1903 Perner: Text-figs 41a, b, p. 67

1963b Horný: Pl. 9, figs 5, 6 (*Bucanella bohemica*)

OUK: Králov Dvůr F.; Lejškov (= Chodouň)

■L 5622, lectotype (SD Horný 1963b)

G = *Tritonophon?* *bohemicus* (PERNER, 1903) (Peel 1974)

bohemica, Carinariopsis: PERNER, 1903

1903 Perner: Pl. 88, figs 41, 42; Text-figs 66a-d, p. 94

1963b Horný: Pl. 10, fig. 1 (*Grandostoma bohemicum*)

1996d Horný: Pl. 2, fig. 4 (*Grandostoma bohemicum*)

OUN: Zahořany F.; Loděnice

■L 5652, holotype by monotypy

G = *Grandostoma bohemicum* (PERNER, 1903)

bohemica, Eobucania?: HORNÝ, 1997

1997c Horný: Text-figs 12c, p. 230; 13a-d, p. 231

OMD: Dobrotivá F.; Praha – Šárka

■L 32361 ±, holotype

G = *Eobucania?* *bohemica* HORNÝ, 1997

bohemica, Kornoutella: (HORNÝ, 1963)

1997c Horný: Text-fig. 20a, p. 235

OML: Šárka F.; Praha – Šárka

L 31986

G? = *Kornoutella bohemica* (HORNÝ, 1963) (Horný 1997c)

bohemicum, Grandostoma: (PERNER, 1903)

1963b Horný: Pl. 10, fig. 4

OUN: Zahořany F.; Praha – Strašnice

L 5656 ±

G = *Grandostoma bohemicum* (PERNER, 1903) (Horný 1963b)

bohemicum, Grandostoma: (PERNER, 1903)

1963b Horný: Pl. 10, figs 5, 6

OUN: Zahořany F.; Loděnice

L 5659

G = *Grandostoma bohemicum* (PERNER, 1903) (Horný 1963b)

- bohemicum, Grandostoma:** (PERNER, 1903)
 1996d Horný: Pl. 1 figs 1, 2; Pl. 2, figs 1-3
 1997b Horný: Text-fig. 1, p. 160 (*Grandostoma bohemicum*)
 OUB: Bohdalec F.; Praha – Hloubětín
L 31170
G = *Grandostoma bohemicum* (PERNER, 1903)
- bohemicum, Grandostoma:** (PERNER, 1903)
 1996d Horný: Pl. 1, figs 3-7; Pl. 2, figs 5, 6; Text-figs 1a, b, p. 224 (non L 31168)
 OUB: Zahořany F.; Praha – Michle
L 31167
G = *Grandostoma bohemicum* (PERNER, 1903)
- bohemicum, Grandostoma:** (PERNER, 1903)
 1996d Horný: Pl. 2, fig. 7; Text-fig. 3, p. 226
 1997b Horný: Text-figs 5a, b, p. 163; Text-fig. 8c, p. 164 (*Grandostoma bohemicum*)
 OUB: Bohdalec F.; Praha – Hloubětín
L 31173
G = *Grandostoma bohemicum* (PERNER, 1903)
- bohemicum, Grandostoma:** (PERNER, 1903)
 1996d Horný: Pl. 2, figs 8, 9
 1997a Horný: Text-fig. 1a, p. 6 (*Grandostoma bohemicum*)
 1997b Horný: Text-figs 4a, b, p. 163; Text-fig. 8b, p. 164 (*Grandostoma bohemicum*)
 OUB: Zahořany F.; Praha – Michle
L 31168
G = *Grandostoma bohemicum* (PERNER, 1903)
- bohemicum, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997b Horný: Text-figs 2a, b, p. 161; Text-figs 3a-c, p. 162; Text-fig. 8a, p. 164 (non L 31168)
 OUB: Zahořany F.; Praha – Spořilov
L 31948
G = *Grandostoma bohemicum* (BARRANDE in PERNER, 1903)
- bohemicum, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997b Horný: Text-fig. 6, p. 163; Text-fig. 8d, p. 134
 OUB: Zahořany F.; Praha – Dubec
L 31951
G = *Grandostoma bohemicum* (BARRANDE in PERNER, 1903)
- bohemicum, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997b Horný: Text-fig. 7, p. 164; text-fig. 8e, p. 164
 OUB: Bohdalec F.; Praha – Hloubětín
L 31952
G = *Grandostoma bohemicum* (BARRANDE in PERNER, 1903)
- bohemicum, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997e Horný: Pl. 1, figs 5, 6
 OUB: Bohdalec F.; Praha – Hloubětín
L 31958
G = *Grandostoma bohemicum* (BARRANDE in PERNER, 1903)
- bohemicum, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997e Horný: Pl. 1, figs 7
 OUB: Bohdalec F.; Praha – Hloubětín
L 31950
G = *Grandostoma bohemicum* (BARRANDE in PERNER, 1903)
- bohemicus, Hypseloconus?**: HORNÝ, 1963
 1963a Horný: Pl. 9, figs 6-10; text-fig. 8, p. 34
 1997c Horný: Text-figs 20b, c, p. 235 (*Kornoutella bohemica*)
 OML: Šárka F.; Praha – Šárka
■L 31985, holotype
G? = *Kornoutella bohemica* (HORNÝ, 1963)
- bouceki, Tropidodiscus:** HORNÝ, 1997
 1997c Horný: Text-fig. 17a, p. 233
 OML: Šárka F.; Praha – Šárka
■L 32380 ±, holotype
G = *Tropidodiscus bouceki* HORNÝ, 1997
- bouceki, Tropidodiscus:** HORNÝ, 1997
 1997c Horný: Text-fig. 17b, p. 233
 OML: Šárka F.; Praha – Šárka
L 32381, paratype
G = *Tropidodiscus bouceki* HORNÝ, 1997
- bouceki, Tropidodiscus:** HORNÝ, 1997
 1997c Horný: p. 233
 OML: Šárka F.; Praha – Šárka
L 32382 ±, paratype
G = *Tropidodiscus bouceki* HORNÝ, 1997
- bouceki, Tropidodiscus:** HORNÝ, 1997
 1997c Horný: p. 233
 OML: Šárka F.; Praha – Šárka
L 32383, paratype
G = *Tropidodiscus bouceki* HORNÝ, 1997
- bouceki, Tropidodiscus:** HORNÝ, 1997
 1997c Horný: p. 233
 OML: Šárka F.; Praha – Šárka
L 32384, paratype
G = *Tropidodiscus bouceki* HORNÝ, 1997
- bouceki, Tropidodiscus:** HORNÝ, 1997
 1997c Horný: p. 233
 OML: Šárka F.; Praha – Šárka
L 32385, paratype
G = *Tropidodiscus bouceki* HORNÝ, 1997
- Bucanella?** sp.
 1997c Horný: Text-fig. 14, p. 231
 1997f Horný: Text-figs 1a-c, p. 334; text-fig. 2a-c, p. 335 (*Tritonophon peeli*)
 1997g Horný: Pl. 8, fig. 7 (*Tritonophon peeli*)
 OUB: Letná F.; Praha – Vysoké Mýto
■L 31999, holotype of *T. peeli*
G = *Tritonophon peeli* HORNÝ, 1997
- Bucanopsis** sp.
 1963b Horný: Pl. 31, fig. 9
 OUK: Králův Dvůr F.; Libomyšl
L 5740
G = *Bucanopsina* sp. (Horný herein)
- Bucanopsis** sp.
 1963b Horný: specimen mentioned on p. 116
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr, Kosov
L 5741
G = *Bucanopsina* sp. (Horný herein)
- calypso, Bucanopsis:** (PERNER, 1903)
 1997a Horný: Text-fig. 1b, p. 6
 OUB: Zahořany F.; Praha – Jinonice
L 31181
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsis:** (PERNER, 1903)
 1997a Horný: Text-figs 5a-d p. 8
 1997b Horný: Text-figs 9a-d, p. 165; Text-fig. 12a, p. 166 (*Bucanopsina calypso*)
 OUB: Zahořany F.; Praha – Michle
L 31174
G = *Bucanopsina calypso* (PERNER, 1903)

- calypso, Bucanopsina:** (PERNER, 1903)
 1997a Horný: Text-figs 6a-c, p. 9; text-figs 7a-c, p. 10
 OUB: Zahořany F.; Praha – Jinonice
L 31180
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997a Horný: Text-fig. 6d, p. 9
 OUB: Zahořany F.; Praha – Libeň, Palmovka
L 31183
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997a Horný: Text-fig. 6e, p. 9
 OUB: Zahořany F.; Praha – Štěrboholy
L 31187
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997a Horný: Text-figs 8a-c, p. 10
 OUB: Zahořany F.; Praha – Jinonice
L 3119
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997a Horný: Text-fig. 9, p. 11
 OUB: Zahořany F.; Praha – Štěrboholy
L 31185
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997a Horný: Text-fig 10 (a), p. 12 (error: *Bucanopsina bohemica*)
 OUB: Letná F.; Revnice, Hvíždinec
L 31945, L 31946, L 31947
G = ? *Bucanopsina roemerii* (BARRANDE in PERNER, 1903) (Horný herein)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997b Horný: Text-fig. 10, p. 165; Text-fig. 12b, p. 166
 OUB: Zahořany F.; Praha – Michle
L 31176
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsina:** (PERNER, 1903)
 1997b Horný: Text-figs 11a, b, p. 166; Text-fig. 12C, p. 166
 OUB: Zahořany F.; Praha – ?Štěrboholy
L 31175
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsis:** PERNER, 1903
 1903 Perner: Text-figs 110a-c, p. 157
 1963b Horný: Pl. 28, figs 6-8 (*Bucanopsis calypso*)
 1997a Horný: Text-figs. 3a-d, p. 7; Text-fig. 4, p. 8 (*Bucanopsis calypso*)
 OUB: Zahořany F.; Praha – Libeň
■L 5738, lectotype (SD Horný 1963b)
G = *Bucanopsina calypso* (PERNER, 1903)
- calypso, Bucanopsis:** PERNER, 1903
 1903 Perner: AD Text-figs 110a-c, p. 157
 1963b Horný: Pl. 28, fig. 9 (*Bucanopsis calypso*)
 OUB: Zahořany F.; Praha – Libeň
L 5739, paralectotype
G = *Bucanopsina calypso* (PERNER, 1903) (Horný 1997a)
- capuloidea, Archinacella:** PERNER, 1903
 1903 Perner: Pl. 1, figs 40, 41; Text-figs 8a-c p. 36
 1963a Horný: Pl. 7, figs 1-5; Text-figs. 6, 7, p. 32 (*Archinacellina modesta*); error: *Archinacella capuloidea*
- 1999 Peel and Horný: Text-figs 10a-f, p. 110 (*Archinacellina modesta*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
■L 5903, holotype by monotypy; by error designated as a paratype of *Archinacella modesta* by Horný 1963a and Peel and Horný 1999
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)
- Carcassonnella** sp. n.: HORNÝ, 1997
 1997c Horný: Text-figs 7a, b, p. 228
 OMD: Dobrotivá F.; Kařízek
L 32400
T = *Carcassonnella* sp. n.
- catenularia, Carinariopsis:** PERNER, 1903
 1903 Perner: Pl. 85, fig. 16
 OUB: Zahořany F.; Loděnice
L 10838, paralectotype
G = *Grandostoma bohemicum* (PERNER, 1903) (Horný 1963b)
- catenularia, Carinariopsis:** PERNER, 1903
 1903 Perner: Pl. 88, fig. 43; Text-figs 65a-d, p. 93
 1963b Horný: Pl. 10, fig. 2 (*Grandostoma bohemicum*)
 OUB: Zahořany F.; Loděnice
■L 5654, lectotype (SD Horný 1996d)
G = *Grandostoma bohemicum* (PERNER, 1903) (Horný 1963b)
- catenularia, Carinariopsis:** PERNER, 1903
 1903 Perner: Pl. 88, fig. 44; Text-fig. 65e, p. 93
 OUB: Zahořany F.; Praha – Štěrboholy
L 5655, paralectotype
G = *Grandostoma bohemicum* PERNER, 1903 (Horný 1963b)
- comata, Bucania:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, figs 30, 31
 1963b Horný: Pl. 29, fig. 6 (*Bucanopsis comata*)
 OUB: Zahořany F.; Zahořany
■L 5742, holotype by monotypy
G = *Bucanopsis cf. calypso* PERNER, 1903 (Horný herein)
- comata, cf., Bucanopsis:** (BARRANDE in PERNER, 1903)
 1963b Horný: Pl. 29, fig. 7
 OUB: Zahořany or Letná Fs.; Vráž
L 5743
G = *Bucanopsis cf. calypso* (BARRANDE in PERNER, 1903) (Horný herein)
- consobrina, Pterotheca:** BARRANDE, 1872
 1872 Barrande: Pl. 20, fig. 19
 1963b Horný: Pl. 16, fig. 4 (*Pterotheca consobrina*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5750, paralectotype
G = *Pterotheca consobrina* BARRANDE, 1872
- consobrina, Pterotheca:** BARRANDE, 1872
 1872 Barrande: Pl. 20, fig. 20
 1963b Horný: Pl. 16, fig. 1 (*Pterotheca consobrina*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
■L 5747, lectotype (SD Horný 1963b)
G = *Pterotheca consobrina* BARRANDE, 1872
- consobrina, Pterotheca:** BARRANDE, 1872
 1872 Barrande: Pl. 20, fig. 21
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5751, paralectotype
G = *Pterotheca consobrina* BARRANDE, 1872

- consobrina, Pterotheca:** BARRANDE, 1872
 1872 Barrande: Pl. 32, fig. 22
 1963b Horný: Pl. 16, fig. 3 (*Pterotheca consobrina*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 L 5749, paralectotype
 G = *Pterotheca consobrina* BARRANDE, 1872
- consobrina, Pterotheca:** BARRANDE, 1872
 1872 Barrande: Pl. 32, fig. 23
 1963b Horný: Pl. 16, fig. 2 (*Pterotheca consobrina*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 L 5748, paralectotype
 G = *Pterotheca consobrina* BARRANDE, 1872
- consobrina, Pterotheca:** BARRANDE, 1872
 1963b Horný: Pl. 16, fig. 5
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 L 5752
 G = *Pterotheca consobrina* BARRANDE, 1872
- Cymbularia** sp.
 1997c Horný: Text-figs 16a, b, p. 233
 OML/OMD: Šárka or Dobrotivá Fs; Praha – Hlubočepy
 L 32397
 G = *Cymbularia* sp.
- delicatula, Bucania:** PERNER, 1903
 1903 Perner: Text-figs 109a-d, p. 156
 1963b Horný: Pl. 10, fig. 3; Pl. 11, fig. 2 (*Grandostoma bohemicum*)
 OUB: Zahořany F.; Loděnice
 ■L 5657, lectotype (SD Horný herein)
 G = *Grandostoma bohemicum* (PERNER, 1903)
- delicatula, Bucania:** PERNER, 1903
 1903 Perner: AD Text-figs 109a-d, p. 156
 OUB: Zahořany F.; Loděnice
 L 5658, paralectotype
 G = *Grandostoma bohemicum* (PERNER, 1903) (Horný herein)
- draboviensis, Bellerophon** (s. l.) (*Bucania*?): BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, fig. 26
 1963b Horný: p. 144 (*species incerta*)
 OUB: Letná F.; Drabov (= Beroun, Děd)
 ■L 10855, holotype by monotypy
 = indeterminate bellerophontiform mollusc
- elevata, Barrandicellosis:** (BARRANDE in PERNER, 1903)
 2000a Horný: Text-fig. 7, p. 149
 OML: Šárka F.; Kařez – Pětidomky
 L 5723
 G = *Barrandicellosis elevata* (BARRANDE in PERNER, 1903)
- elevata, Barrandicellosis:** (BARRANDE in PERNER, 1903)
 2000a Horný: Text-figs 8a, b, p. 149
 OML: Šárka F.; Osek
 L 30063
 G = *Barrandicellosis elevata* (BARRANDE in PERNER, 1903)
- evolvens, Cyrtolites?**: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 86, figs 22-24
 1963b Horný: Pl. 15, fig. 2 (*Temnodiscus evolvens*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr, Kosov
 ■L 5670, holotype by monotypy
 T = *Cyrtodiscus?* *evolvens* (BARRANDE in PERNER, 1903) (Horný herein)
- evolvens, Temnodiscus:** (BARRANDE in PERNER, 1903)
 1963b Horný: Pl. 15, fig. 1
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr, Kosov
 L 5671
 T = *Cyrtodiscus?* *evolvens* (BARRANDE in PERNER, 1903) (Horný herein)
- extenuata, Orthonychia:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 3, figs 11, 12
 1911 Perner: p. 186 (*Orthonychia?* *extenuata*)
 2000a Horný: Text-figs 11Aa, b, p. 151 (*Barrandicellosis?* *extenuata*)
 OUB: Letná F.; Drabov (= Beroun, Děd)
 ■L 5425, holotype by monotypy
 G = *Barrandicellosis?* *extenuata* (BARRANDE in PERNER, 1903)
- extenuata, Barrandicellosis?:** (BARRANDE in PERNER, 1903)
 2000a Horný: Text-figs 11Ba, b, p. 151
 OUB: Letná F.; fields between Trubská and Trubín
 L 32742
 G = *Barrandicellosis?* *extenuata* (BARRANDE in PERNER, 1903)
- extenuata, Barrandicellosis?:** (BARRANDE in PERNER, 1903)
 2000a Horný: Text-figs 11Ca, b, p. 151
 OUB: Letná F.; fields between Trubská and Trubín
 L 32743
 G = *Barrandicellosis?* *extenuata* (BARRANDE in PERNER, 1903)
- extenuata, Barrandicellosis?:** (BARRANDE in PERNER, 1903)
 2000a Horný: Text-fig. 11D, p. 151
 OUB: Letná F.; fields between Trubská and Trubín
 L 32744
 G = *Barrandicellosis?* *extenuata* (BARRANDE in PERNER, 1903)
- ferrigena, Temnodiscus:** PERNER, 1903
 1903 Perner: Text-fig. 54, p. 78
 1907 Perner: Pl. 114, figs 5, 6 (*Sinuitopsis nodosa?*)
 OUB: Zahořany F.; Loděnice
 ■L 5630, holotype by monotypy
 T = *Sinuitopsis neglecta* PERNER, 1903 (Horný 1963b)
- grande, Grandostoma:** (BARRANDE in PERNER, 1903)
 1963b Horný: Pl. 11, fig. 3
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 L 5663
 G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande, Grandostoma:** (BARRANDE in PERNER, 1903)
 1963b Horný: specimen mentioned on p. 86 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
 L 5665
 G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997e Horný: Pl. 1, fig. 1
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
 L 31960
 G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande, Grandostoma:** (BARRANDE in PERNER, 1903)
 1997e Horný: Pl. 1, fig. 2
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
 L 31961
 G = *Grandostoma grande* (PERNER, 1903)
- grande, Salpingostoma:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 84, figs 12-14
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 L 36540, paralectotype
 G = *Grandostoma grande* (BARRANDE in PERNER, 1903) (Horný 1963b)
- grande, Salpingostoma:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, figs 25
 1963b Horný: Pl. 11, fig. 4 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 L 5662, paralectotype
 G = *Grandostoma grande* (BARRANDE in PERNER, 1903)

- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, figs 26, 27
 1963b Horný: Pl. 12, fig. 3 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5666, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, fig. 28
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 10840, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903) (Horný 1963b)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, fig. 29
 1963b Horný: Pl. 12, figs 5, 6 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5661, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, fig. 30; Text-fig. 69f, p. 98
 1963b Horný: Pl. 12, fig. 2 (*Grandostoma bohemicum*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
■L 5660, lectotype (SD Horný 1963b)
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, fig. 31
 1963b Horný: Pl. 11, fig. 5; Pl. 12, fig. 1 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 5667, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, fig. 32
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 36709, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903) (Horný 1963b)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Text-figs 69a₁, a₂, p. 98
 1997c Horný: Pl. 1, figs 3, 4 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 5668, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Text-figs 69b₁, b₂, p. 98
 1963b Horný: Pl. 1, fig. 6 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 5669, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Text-fig. 69e, p. 98
 1963b Horný: Pl. 12, fig. 4 (*Grandostoma grande*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 5664, paralectotype
G = *Grandostoma grande* (BARRANDE in PERNER, 1903)
- grande?*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, fig. 1
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 10852
G = *Sinuitopsis aff. bilobatus* (SOWERBY, 1839) (Horný herein)
- grande?*, *Salpingostoma*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, fig. 2
 OUK: Králův Dvůr F., Beroun – Králův Dvůr
L 10851
G = *Bucanopsisina?* sp. (Horný herein)
- hanusi*, *Sinuites*: HORNY, 1997
 1997c Horný: Text-figs 9a-c, p. 227
 OML: Šárka F.; Praha – Šárka
■L 32356 ±, holotype
G = *Sinuites hanusi* HORNY, 1997
- hanusi*, *Sinuites*: HORNY, 1997
 1997c Horný: p. 229
 OML: Šárka F.; Praha – Šárka
L 32357, paratype
G = *Sinuites hanusi* HORNY, 1997
- hanusi*, *Sinuites*: HORNY, 1997
 1997c Horný: p. 229
 OML: Šárka F.; Praha – Šárka
L 32358 ±, paratype
G = *Sinuites hanusi* HORNY, 1997
- hanusi*, *Sinuites*: HORNY, 1997
 1997c Horný: p. 229
 OML: Šárka F.; Praha – Šárka
L 32359 ±, paratype
G = *Sinuites hanusi* HORNY, 1997
- hanusi*, *Sinuites*: HORNY, 1997
 1997c Horný: p. 229
 OML: Šárka F.; Praha – Šárka
L 32360 ±, paratype
G = *Sinuites hanusi* HORNY, 1997
- hanusi*, *Sinuites*: HORNY, 1997
 1997c Horný: specimen mentioned on p. 229
 OML: Šárka F.; Praha – Šárka
L 32401
G = *Sinuites hanusi* HORNY, 1997
- immigrans*, *Palaeacmaea*: BARRANDE in PERNER, 1903
 1907 Perner: Pl. 104, figs 32-34
 1963a Horný: Pl. 10, fig. 7 (*Patelliconus primulus*)
 OML: Šárka F.; Osek
■L 11685, holotype by monotypy; by error designated as a paratype of *Palaeacmaea primula* by Horný 1963a
G = *Patelliconus primulus* (BARRANDE in PERNER, 1903)
- incertus*, *Temnодiscus*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, figs 12, 13; Text-figs 53a, b, p. 78
 OUB: Letná F.; Trubská
■L 5645, holotype by monotypy
T = *Sinuitopsis neglecta* (BARRANDE in PERNER, 1903) (Horný 1963b)
- incertus*, cfr., *Temnодiscus*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 42, figs 35, 36
 OUB: Letná F.; Drabov (= Beroun, Děd)
L 8679
T = *Sinuitopsis neglecta* (BARRANDE in PERNER, 1903) (Horný herein)
- incertus?*, *Temnодiscus*: PERNER, 1903
 1903 Perner: Pl. 42, fig. 37
 OUB: Letná F.; Trubská
L 8680
T = *Sinuitopsis neglecta* (BARRANDE in PERNER, 1903) (Horný herein)

- incola, Bellerophon* (s. l.) (*Oxydiscus?*): BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, figs 9-11
 1963b Horný: p. 144 (*species incerta*)
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
 ■L 10854, holotype by monotypy
 G = ? *Simites* sp. (specimen damaged)
- kettneri, Palaeacmaea*: ŘÍHA, 1938
 1938 Říha: Pl. I, figs 2a, b 1963a
 1963a Horný: Pl. 9, fig. 11 (*Pygmaeconus? kettneri*)
 OML: Šárka F.; Osek
 ■L 30070, holotype by monotypy
 G = *Pygmaeconus kettneri* (ŘÍHA, 1938) (Horný herein)
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: Text-fig. 15, p. 232
 OMD: Dobrotivá F.; Malé Přílepy
 ■L 32386, holotype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: Text-fig. 15, p. 232
 OMD: Dobrotivá F.; Malé Přílepy
 L 32387, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Malé Přílepy
 L 32388, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32389, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32390, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32391, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32392, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32393, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32394, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32395, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- klouceki, Cymbularia*: HORNÝ, 1997
 1997c Horný: p. 232
 OMD: Dobrotivá F.; Praha – Šárka, a field at the villa
 L 32396, paratype
 G = *Cymbularia klouceki* HORNÝ, 1997
- kolebabai, Sarkanella*: HORNÝ, 2002
 2002a Horný: Text-fig. 1, in press
 OUK: Králův Dvůr F.; Praha – Pankrác
 ■L 32751, holotype
 T = *Sarkanella kolebabai* HORNÝ, 2002
- krafti, Pygmaeconus*: HORNÝ, 1997
 1997c Horný: Text-fig. 21a, b, p. 235
 OUB: Zahořany F.; Praha – Libeň, railway cut
 ■L 7678, holotype
 G = *Pygmaeconus krafti* HORNÝ, 1997
- krafti, Pygmaeconus*: HORNÝ, 1997
 1997c Horný: p. 235
 OUB: Bohdalec F.; Praha – Michle, a shunting yard
 L 5910, paratype
 G = *Pygmaeconus krafti* HORNÝ, 1997
- krafti, Pygmaeconus*: HORNÝ, 1997
 1997c Horný: p. 235
 OUB: Zahořany F.; Praha – Nový Hloubětín
 L 7679 ±, paratype
 G = *Pygmaeconus krafti* HORNÝ, 1997
- krafti, Pygmaeconus*: HORNÝ, 1997
 1997c Horný: p. 235
 OUB: Zahořany F.; Praha – between Dubeč and Horní Počernice
 L 31988, paratype
 G = *Pygmaeconus krafti* HORNÝ, 1997
- krafti, Pygmaeconus*: HORNÝ, 1997
 1997c Horný: p. 235
 OUB: Zahořany F.; Loděnice, N of the Kněží hora Hill
 L 31989 ±, paratype
 G = *Pygmaeconus krafti* HORNÝ, 1997
- latiuscula, Palaeacmaea*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 5, figs 18, 19
 1963a Horný: Pl. 18, fig. 6 (*species dubia*)
 OML: Šárka F.; Osek
 ■L 8386, holotype by monotypy
 G = *Pygmaeconus? latiusculus* (BARRANDE in PERNER, 1903)
- mergli, Peelerophon*: (FRÝDA, 1988)
 1997c Horný: Text-figs 5a-c, p. 227
 OLA: Klabava F.; Kleštěnice near Komárov
 L 31997
 T = *Peelerophon mergli* (FRÝDA, 1988)
- mitra, Archinacella*: PERNER, 1903
 1903 Perner: Pl. I, figs 38, 39
 1963b Horný: Pl. 18, figs 4, 5 (? *Archinacellina modesta*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 ■L 5882, holotype by monotypy
 G = *Archinacellina modesta* (BARRANDE in PERNER, 1903) (Horný herein)

- modesta, Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 1, figs 25, 26
 1963a Horný: Pl. 6, figs 10, 11 (*Archinacellina modesta*)
 OUK: Králov Dvůr F.; Lejškov (= Chodouň)
L 5862, paralectotype
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)
- modesta, Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 1, figs 27, 28
 1963a Horný: Pl. 6, fig. 9 (*Archinacellina modesta*)
 OUK: Králov Dvůr F.; Lejškov (= Chodouň)
L 5872, paralectotype
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)
- modesta, Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 1, figs 42, 43
 1963a Horný: Pl. 8, fig. 3 (*Archinacellina modesta*)
 OUK: Králov Dvůr F.; Lejškov (= Chodouň)
L 5865, paralectotype
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)
- modesta, Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 1, figs 44, 45
 1963a Horný: Pl. 8, figs 1, 2 (*Archinacellina modesta*)
 OUK: Králov Dvůr F.; Lejškov (= Chodouň)
■L 5864, lectotype (SD Horný 1963a)
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)
- modesta?, Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 1, figs 29, 30
 1963a Horný: Pl. 8, fig. 4 (not 5) (? *Archinacella tarda*)
 OUB: Zahořany F.; Loděnice
L 5879
G = ? *Barrandicella?* *tarda* (PERNER, 1903) (Horný herein)
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 85, fig. 11
 1991b Horný: Pl. 10, figs 3-5 (*Sinuitopsis neglecta*)
 OUB: Zahořany F.; Praha – Libeň
L 10837, several paralectotypes
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 88, figs 28-30
 OUB: Zahořany F.; Loděnice
L 10862, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1903 Perner: Text-figs 42a-f, p. 69; Text-fig. 44a, p. 70
 1963b Horný: Pl. 9, fig. 4 (*Sinuitopsis neglecta*)
 1990 Horný: Pl. 1, fig. 5 (*Sinuitopsis neglecta*)
 1991b Horný: Pl. 1, figs 8-10; Text-fig. 5/6, p. 87 (*Sinuitopsis neglecta*)
 OUB: Zahořany F.; Loděnice or Praha – Libeň
L 5625, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1903 Perner: Text-fig. 43a-e p. 70
 1941 Knight: Pl. 6, figs 4a-c (*Sinuitopsis neglecta*)
 1963b Horný: Pl. 6, figs 5, 6; Pl. 7, fig. 3 (*Sinuitopsis neglecta*)
 1990 Horný: Pl. 1, fig. 6 (*Sinuitopsis neglecta*)
 1991b Horný: Pl. 1, figs 1-7; Text-fig. 5/3, p. 87 (*Sinuitopsis neglecta*)
 OUB: Zahořany F.; Loděnice
■L 5624, lectotype (SD Knight 1941)
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1903 Perner: Text-figs 44b, d, p. 70
 OUB: Zahořany F.; Loděnice
L 36536, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1903 Perner: Text-figs 44c, p. 70
 OUB: Zahořany F.; Loděnice
L 36537, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1963b Horný: specimen mentioned on p. 78 (*Sinuitopsis neglecta*)
 OML: Šárka F.; Praha – Úvaly
L 5631
T = *Sinuitopsis cf. evoluta* (BARRANDE in PERNER, 1903) (Horný herein)
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1963b Horný: specimen mentioned on p. 78 (*Sinuitopsis neglecta*)
 OML: Šárka F.; Praha – Úvaly
L 5632
T = *Sinuitopsis cf. evoluta* (BARRANDE in PERNER, 1903) (Horný herein)
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 2, figs 1, 2; Pl. 9, fig. 1
 OUB: Zahořany F.; Praha – Vysočany
L 28629 ±
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 2, figs 3-9; Text-fig. 11, p. 91
 OUB: Zahořany F.; Praha – Libeň
L 28630
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 3, figs 1-6; Text-fig. 1, p. 83, Text-fig. 4, p. 86
 (partim); Text-fig. 5/9, p. 87
 OUB: Zahořany F.; Praha – Spořilov
L 28631
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 3, fig. 7
 OUB: Zahořany F.; Praha – Jinonice
L 28632
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 4, figs 1-3; Text-fig. 5/2, p. 87; Text-fig. 9 (left), p. 89
 1991 Peel: Text-figs 18d-f, p. 24 (*Sinuitopsis neglecta*)
 OUB: Zahořany F.; Loděnice or Praha – Libeň
L 28633
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 4, figs 4-6; Text-fig. 5/8, p. 87; Text-fig. 9 (right), p. 89
 OUB: Zahořany F.; ? Praha – Radotín
L 28634
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- neglecta, Sinuitopsis:** BARRANDE in PERNER, 1903
 1991b Horný: Pl. 4, figs 7, 8; Text-fig. 5/4, p. 87
 OUB: Zahořany F.; Praha – Libeň
L 28635
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 4, fig. 9; Pl. 5, figs 1-8; Text-fig. 5/5, p. 87
OUB: Zahořany F.; Praha – Spořilov
L 28636 ±
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 6, figs 1-6; Pl. 7, figs 1-9; Text-fig. 5/7, p. 87
OUB: Zahořany F.; Praha – Vysočany
L 28637 ±
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 8, fig. 1; Text-fig. 12, p. 92
OUB: Zahořany F.; Praha – Libeň
L 28638
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 9, fig. 2
OUB: Zahořany F.; Praha – Krč
L 28639
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 9, fig. 3
OUB: Zahořany F.; Praha – Libeň
L 28895
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 9, fig. 4
OUB: Zahořany F.; Praha – Libeň or Loděnice
L 28896
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 10, fig. 7
OUB: Zahořany F.; Praha – Vysočany
L 28641 ±
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 12, figs 1-3, Text-fig. 5/1, p. 87
OUB: Zahořany F.; Praha – Vysočany
L 28642
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 12, fig. 4; Text-fig. 8 (left), p. 88
OUB: Zahořany F.; Praha – Jinonice
L 28643
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 12, fig. 5
OUB: Zahořany F.; Praha – Jinonice
L 28644
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 12, fig. 6
OUB: Zahořany F.; Praha – Jinonice
L 28897
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Pl. 12, fig. 7; Text-fig. 8 (right), p. 88
OUB: Zahořany F.; Praha – Jinonice
L 28898
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: AD Text-fig. 4, p. 86
OUB: Zahořany F.; Praha – Spořilov
L 28900
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: AD Text-fig. 12, p. 92
OUB: Zahořany F.; Praha – Libeň
L 28901
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1991b Horný: Text-fig. 13, p. 93
OUB: Zahořany F.; Praha – Libeň
L 28899
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1997a Horný: Text-fig. 10 (b), p. 12
OUB: Letná F.; Řevnice, Hvíždinec
L 31947
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: BARRANDE in PERNER, 1903
1997d Horný: Pl. 1, figs 1-3
OUB: Zahořany F.; Praha – Jinonice
L 31955
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta, *Sinuitopsis*: PERNER, 1903
1997d Horný: Pl. 1, figs 4-6; Pl. 2, figs 1, 2; Pl. 3, figs 1-3, Pl. 4, figs 1-3
OUB: Zahořany F.; Praha – Spořilov
L 31954
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta var. *transgrediens*, *Sinuitopsis*: PERNER, 1903
1903 Perner: Pl. 88, fig. 33
1963b Horný: Pl. 7, figs 4-6 (*Sinuitopsis neglecta*)
OUB: Zahořany F.; ? Loděnice
L 5627, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta var. *transgrediens*, *Sinuitopsis*: PERNER, 1903
1903 Perner: Pl. 88, figs 35-37
1963b Horný: Pl. 7, figs 7, 8 (*Sinuitopsis neglecta*)
OUB: Zahořany F.; Praha – Radotín
L 5628, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903

neglecta var. *transgrediens*, *Sinuitopsis*: PERNER, 1903
1903 Perner: Text-figs 45a, b, p. 71
OUB: Zahořany F.; Butovice (= Praha – Jinonice)
L 23771, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný 1963b)

neglecta var. *transgrediens*, *Sinuitopsis*: PERNER, 1903
1903 Perner: Text-figs 46a, b, p. 71
OUB: Zahořany F.; Praha – Libeň
L 10863, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný 1963b)

neglecta var. *transgrediens*, *Sinuitopsis*: PERNER, 1903
1903 Perner: Text-fig. 46c, p. 71
1963b Horný: Pl. 7, fig. 2 (*Sinuitopsis neglecta*)

- OUB: Zahořany F; Praha – Štěrbohol
L 5626, lectotype (SD Horný herein)
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903
- nitidus**, *Oxydiscus* (*Cyrtodiscus*): BARRANDE in PERNER, 1903
1903 Perner: Pl. 86, figs 8-13
1963b Horný: Pl. 13, figs 3, 4 (*Gamadiscus nitidus*)
OML: Šárka F.; Osek
L 5830, lectotype (SD Horný 1963b)
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Oxydiscus* (*Cyrtodiscus*): BARRANDE in PERNER, 1903
1903 Perner: AD Pl. 86, fig. 8
1963b Horný: Pl. 13, fig. 3 (*Gamadiscus nitidus*)
OML: Šárka F.; Osek
L 5830, 3 paralectotypes
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Oxydiscus* (*Cyrtodiscus*): BARRANDE in PERNER, 1903
1903 Perner: AD Pl. 86, fig. 8
OML: Šárka F.; Osek
L 5829, several paralectotypes
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Gamadiscus*: (BARRANDE in PERNER, 1903)
1963b Horný: Pl. 13, fig. 5
1996 Horný and Peel: Pl. 7, fig. 7 (*Gamadiscus nitidus*)
OML: Šárka F.; Osek
L 5831
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Gamadiscus*: (BARRANDE in PERNER, 1903)
1963b Horný: Pl. 13, fig. 6
1996 Horný and Peel: Pl. 7, fig. 8 (*Gamadiscus nitidus*)
OML: Šárka F.; Osek
L 5832
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Gamadiscus*: (BARRANDE in PERNER, 1903)
1963b Horný: Pl. 13, fig. 7
OML: Šárka F.; Osek
L 5833
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Gamadiscus*: (BARRANDE in PERNER, 1903)
1963b Horný: Pl. 13, fig. 8
OML: Šárka F.; Osek
L 5834
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Gamadiscus*: (BARRANDE in PERNER, 1903)
1996b Horný: Text-figs 1a, b, p. 245; Text-fig. 4a, p. 247
OML: Šárka F.; Osek
L 31754
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Gamadiscus*: (BARRANDE in PERNER, 1903)
1996b Horný: Text-fig. 4c, p. 247
- OML: Šárka F.; Osek
L 31755
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Cyrtodiscus*: (PERNER, 1903)
1998 Horný: Text-figs 4a-d, p. 213
OML: Šárka F.; Osek
L 32549
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Cyrtodiscus*: (PERNER, 1903)
1998 Horný: Text-figs 5a, p. 214; Text-fig. 7f, p. 216
OML: Šárka F.; Kařez – Pětidomky
L 32546
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Cyrtodiscus*: (PERNER, 1903)
1998 Horný: Text-figs 5d, p. 214; Text-fig. 7c, p. 216
OML: Šárka F.; Kařez – Pětidomky
L 32548
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nitidus**, *Cyrtodiscus*: (PERNER, 1903)
1998 Horný: Text-figs 5g, p. 214; Text-fig. 7g, p. 216
OML: Šárka F.; Kařez – Pětidomky
L 32547
T = *Cyrtodiscus nitidus* (BARRANDE in PERNER, 1903) (Horný 1997g)
- nodos**, *Sinuitopsis*: PERNER, 1903
1903 Perner: Pl. 88, figs 31, 32
OUB: Zahořany F.; Loděnice
L 5636, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný herein)
- nodos**, *Sinuitopsis*: PERNER, 1903
1903 Perner: Text-figs 48a-c, p. 72
1963b Horný: Pl. 7, fig. 1 (*Sinuitopsis nodosa*)
OUB: Zahořany F.; Loděnice
L 5633, lectotype (SD Horný 1963b)
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný herein)
- nodos**, *Sinuitopsis*: PERNER, 1903
1903 Perner: Text-fig. 48d, p. 72
OUB: Zahořany F.; Loděnice
L 23776, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný herein)
- nodos**, *Sinuitopsis*: PERNER, 1903
1903 Perner: Text-figs 49a, b, p. 73
1963b Horný: specimen mentioned on p. 79 (*Sinuitopsis nodosa*)
OUB: Zahořany F.; Loděnice
L 5634, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný herein)
- nodos**, *Sinuitopsis*: PERNER, 1903
1903 Perner: AD Text-figs 49a, b, p. 73
1963b Horný: specimen mentioned on p. 79 (*Sinuitopsis nodosa*)
OUB: Zahořany F.; Loděnice
L 5635, paralectotype
T = *Sinuitopsis neglecta* BARRANDE in PERNER, 1903 (Horný herein)

- novaki, Sarkanella:** HORNÝ et VONKA, 2002
 2002 Horný et Vonka: Pl. I, figs 1-3
 OUB: Bohdalec F.; Praha – Radotín
 ■L 32406, holotype
 T = *Sarkanella novaki* HORNÝ et VONKA, 2002
- novaki, Sarkanella:** HORNÝ et VONKA, 2002
 2002 Horný et Vonka: Pl. I, fig. 4
 OUB: Bohdalec F.; Praha – Radotín
 L 32750, paratype
 T = *Sarkanella novaki* HORNÝ et VONKA, 2002
- ordovicina, Palaeoscurria:** HORNÝ, 1963
 1963a Horný: Pl. 17, figs 7-9
 OMD: Dobrotivá F.; Malé Přílepy
 ■L 5857, holotype
 G = *Micropileus?* *ordovicinus* (HORNÝ, 1963) (Horný herein)
- ovata, Archinacella:** BARRANDE in PERNER, 1903
 1963a Horný: Pl. 4, fig. 4
 2000b Horný: Pl. I, fig. 3 (*Barrandicella ovata* + *Barrandicella holubi*)
 OML: Šárka F.; Osek
 L 5875, paralectotypes of *B. ovata* + L 32745, a paratype of *B. holubi*
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Peel and Horný 1999)
 G = *Barrandicella holubi* HORNÝ, 2000
- ovata, Archinacella:** BARRANDE in PERNER, 1903
 1963a Horný: Pl. 5, figs 9, 10
 2000b Horný: Pl. I, figs 8, 9; Pl. 2, fig. 1 (*Barrandicella holubi*)
 OML: Šárka F.; Osek
 ■L 5896, holotype of *Barrandicella holubi*
 G = *Barrandicella holubi* HORNÝ, 2000
- ovata var. complanata, Archinacella:** PERNER, 1903
 1903 Perner: Text-figs 5a, b, p. 33
 1963a Horný: Pl. 6, figs 3-5 (*Archinacella ovata*)
 OML: Šárka F.; Osek
 ■L 16299, lectotype (SD Horný herein)
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Horný herein)
- ovata var. complanata, Archinacella:** PERNER, 1903
 1903 Perner: AD Text-figs 5a, b, p. 33
 1963a Horný: Pl. 4, fig. 5 (*Archinacella ovata*)
 OML: Šárka F.; Osek
 L 16300, paralectotype
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Horný herein)
- ovata var. elevata, Archinacella:** PERNER, 1903
 1903 Perner: p. 33
 1907 Perner: Pl. 104, figs 30, 31 (*Archinacella ovata* var. *elevata*)
 1963a Horný: pp. 26, 28 (*Archinacella?* *ovata elevata*)
 2000a Horný: Text-figs 9a, b, p. 150 (*Barrandicellopsis elevata*)
 OML: Šárka F.; Osek
 ■L 5722, holotype by monotypy
 G = *Barrandicellopsis elevata* (PERNER, 1903) (Horný 2000a)
- ovata var. rostrata, Archinacella:** PERNER, 1903
 1903 Perner: Pl. 2, figs 1, 2
 1963a Horný: Pl. 4, fig. 6 (*Archinacella ovata*)
 OML: Šárka F.; Osek
- L 31751, paralectotype
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Horný 2000a)
- ovata var. rostrata, Archinacella:** PERNER, 1903
 1903 Perner: Text-figs 4a, b, p. 33
 1963a Horný: Pl. 6, figs 6-8 (*Archinacella ovata*)
 2000a Horný: Text-fig. 3, p. 147 (*Barrandicellopsis elevata*)
 OML: Šárka F.; Osek
 ■L 16298, lectotype (SD Horný 2000)
 G = *Barrandicellopsis elevata* (PERNER, 1903)
- ovata var. subglobosa, Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Text-figs 6a, b, p. 34
 1963a Horný: Pl. 7, fig. 8 (*Archinacella ovata*)
 OML: Šárka F.; Osek
 ■L 5863, holotype by monotypy
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Horný herein)
- ovata var. tarda, Archinacella:** PERNER, 1903
 1903 Perner: Pl. 43, figs 22, 23
 1963a Horný: Pl. 6, fig. 2 (*Archinacella tarda*)
 OUB: Zahofány F.; Vráž
 L 5901, paralectotype
 G = *Barrandicella?* *tarda* (PERNER, 1903) (Horný herein)
- ovata var. tarda, Archinacella:** PERNER, 1903
 1903 Perner: Text-figs 7a, b, p. 34
 1963a Horný: Pl. 6, fig. 1; Pl. 7, figs. 6, 7 (*Archinacella tarda*)
 OUB: Zahofány F.; Loděnice
 ■L 5886, lectotype (SD Horný 1963a)
 G = *Barrandicella?* *tarda* (PERNER, 1903) (Horný herein)
- ovata var. typica [= ovata], Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 2, figs 3, 4
 OML: Šárka F.; Osek
 L 5887, paralectotype
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Peel and Horný 1999)
- ovata var. typica [= ovata], Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 2, figs 5, 6
 1963a Horný: Pl. 4, fig. 6 (*Archinacella ovata*)
 OML: Šárka F.; Osek
 L 5866, paralectotype
 G = *Barrandicella?* *ovata* (BARRANDE in PERNER, 1903) (Peel and Horný 1999)
- ovata var. typica [= ovata], Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 2, figs 7, 8; ? Text-figs 3a, b + 3bis
 1963a Horný: Pl. 5, figs 3-5 (*Archinacella ovata*)
 OML: Šárka F.; Osek
 ■L 5861, lectotype (SD Horný 1963a)
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Peel and Horný 1999)
- ovata var. typica [= ovata], Archinacella:** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 2, figs 9, 10
 1963a Horný: Pl. 5, figs 6-8 (*Archinacella ovata*)
 OML: Šárka F.; Osek
 L 8381, paralectotype
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903) (Peel and Horný 1999)

- ovata, Barrandicella*: (BARRANDE in PERNER, 1903)
 1999 Peel and Horný: Text-figs 6a-c, p. 106
 OML: Šárka F.; Osek
L 31757
 G = *Barrandicella ovata* (BARRANDE in PERNER, 1903)
- peeli, Tritonophon*: HORNÝ, 1997
 1997f Horný: Text-figs 4a-c, p. 335
 OUB: Letná F.; Praha – Vysočany
L 32402, paratype
 G = *Tritonophon peeli* HORNÝ, 1997
- perneri, Selesinuites*: HORNÝ, 1997
 1997c Horný: Text-figs 12a, b, p. 230
 1997g Horný: Pl. 7, fig. 12
 OML: Šárka F.; Osek
■L 32362, holotype
 G = *Selesinuites perneri* HORNÝ, 1997
- plasi, Sinuitopsisina*: HORNÝ, 1997
 1997c Horný: Text-figs 3a-c, p. 226
 OMD: Dobrotivá F.; Praha – Šárka
■L 31978, holotype
 T = *Sinuitopsisina plasi*: HORNÝ, 1997
- platynotus, Temnodiscus*: PERNER, 1903
 1903 Perner: Text-figs 51a-c, p. 76
 1963b Horný: p. 77 (*Sinuitopsisina neglecta*)
 OUB: Letná F.; Drabov (= Beroun, Děd)
■L 5639, holotype by monotypy
 T = *Sinuitopsisina neglecta* BARRANDE in PERNER, 1903
- podexpulicis, Marekicella*: HORNÝ, 1997
 1997c Horný: Textfigs 22a-c, p. 236
 OUB: Zahořany F.; Staňkovka near Praha – Radotín
■L 31974 ±, holotype
 G = *Marekicella podexpulicis* HORNÝ, 1997
- podexpulicis, Marekicella*: HORNÝ, 1997
 1997c Horný: p. 236
 OUB: Bohdalec F.; Praha – Vršovice
L 31975 ±, paratype
 G = *Marekicella podexpulicis* HORNÝ, 1997
- orrecta, Palaeacmaea?*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 5, figs 4-6
 OML: Šárka F.; Osek
L 8269, paralectotype
 G = *Pygmaeoconus porrectus* (BARRANDE in PERNER, 1903)
 (Horný 1963a)
- orrecta, Palaeacmaea?*: BARRANDE in PERNER, 1903
 1903 Perner: p. 29
 1911 Perner: Pl. 247, figs 1-3 (*Palaeacmaea?* *orrecta*)
 1961 Horný: Pl. 2, fig. 2 (*Pygmaeoconus porrectus*)
 1963a Horný: Pl. 9, figs 3-5; Text-fig. 12, p. 38 (*Pygmaeoconus porrectus*)
 OML: Šárka F.; Osek
■L 29379, lectotype (SD Horný 1961 as the holotype)
 G = *Pygmaeoconus porrectus* (BARRANDE in PERNER, 1903)
- orrectus, Pygmaeoconus*: (BARRANDE in PERNER, 1903)
 1961 Horný: Pl. 2, fig. 1
 1963a Horný: Pl. 9, fig. 1; Text-fig. 11, p. 38 (*Pygmaeoconus porrectus*)
 OML: Šárka F.; Osek
L 5877, by error designated as a paratype by Horný 1961
 G = *Pygmaeoconus porrectus* (BARRANDE in PERNER, 1903)
- orrectus, Pygmaeoconus*: (BARRANDE in PERNER, 1903)
 1963a Horný: Pl. 9, fig. 2
- OML: Šárka F.; Osek
L 5858
 G = *Pygmaeoconus porrectus* (BARRANDE in PERNER, 1903)
- orrectus, Pygmaeoconus*: (BARRANDE in PERNER, 1903)
 2002b Horný: Pl. 4, fig. 1
 OML: Šárka F., Osek
L 31982
 G = *Pygmaeoconus porrectus* (BARRANDE in PERNER, 1903)
- ragensis, Carcassonnella*: HORNÝ, 1997
 1997c Horný: Text-fig. 6a, p. 227
 OML: Šárka F.; Praha – Šárka
■L 32372, holotype
 T = *Carcassonnella ragensis* HORNÝ, 1997
- ragensis, Carcassonnella*: HORNÝ, 1997
 1997c Horný: Text-fig. 6b, p. 227
 OML: Šárka F.; Praha – Šárka
L 32373 ±, paratype
 T = *Carcassonnella ragensis* HORNÝ, 1997
- ragensis, Carcassonnella*: HORNÝ, 1997
 1997c Horný: Text-fig. 6c, d, p. 227
 OML: Šárka F.; Praha – Šárka
L 32374, paratype
 T = *Carcassonnella ragensis* HORNÝ, 1997
- ragensis, Carcassonnella*: HORNÝ, 1997
 1997c Horný: p. 227
 OML: Šárka F.; Praha – Šárka
L 32375, paratype
 T = *Carcassonnella ragensis* HORNÝ, 1997
- ragensis, Carcassonnella*: HORNÝ, 1997
 1997c Horný: p. 227
 OML: Šárka F.; Praha – Šárka
L 32376, paratype
 T = *Carcassonnella ragensis* HORNÝ, 1997
- rantli, Pentalina*: HORNÝ, 1961
 1961 Horný: Pl. 2, fig. 3
 1963a Horný: Pl. 3, figs 1-5; Text-fig. 1, p. 15
 OMD: Dobrotivá F.; Malé Přílepy
■L 5403, holotype
 T = *Pentalina rantli* HORNÝ, 1961
- rimula, Palaeacmaea*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 48, figs 4-6
 1961 Horný: *Patelliconus primulus*
 1963a Horný: Pl. 10, figs 4-6; Text-fig. 10, p. 36 (*Patelliconus primulus*)
 OML: Šárka F.; Osek
■L 8425, holotype by monotypy
 G = *Patelliconus primulus* (BARRANDE in PERNER, 1903)
- rimulus, Patelliconus*: (BARRANDE in PERNER, 1903)
 1963a Horný: Pl. 10, figs 1-3; Text-fig. 9, p. 36
 OML: Šárka F.; Osek
L 5891
 G = *Patelliconus primulus* (BARRANDE in PERNER, 1903)
- rocer, Oxydiscus (Cyrtodiscus)*: BARRANDE in PERNER, 1903
 1903 Perner: Pl. 88, figs 22-24; Text-figs 50a, b, p. 74
 1941 Knight: Pl. 9, fig. 4a (*Cyrtodiscus procer*)
 1963b Horný: Pl. 14, figs 1, 2 (*Cyrtodiscus procer*)
 OUB: Letná F.; Trubská
■L 5679, lectotype (SD Knight 1941)
 T = *Cyrtodiscus procer* (BARRANDE in PERNER, 1903)

procer, *Oxydiscus (Cyrtodiscus)*: BARRANDE in PERNER, 1903
1903 Perner: AD Pl. 88, figs 22-24; ADText-figs 50a, b, p. 74
1941 Knight: Pl. 9, fig. 4b (*Cyrtodiscus procer*)
1963b Horný: Pl. 14, fig. 3 (*Cyrtodiscus procer*)
OUB: Letná F.; Trubská
L 5680, paralectotype
T = *Cyrtodiscus procer* (BARRANDE in PERNER, 1903)

procer, *Cyrtodiscus*: (BARRANDE in PERNER, 1903)
1997c Horný: Text-fig. 1a, p. 224
OUB: Letná F.; Praha – Vysočany
L 31990
T = *Cyrtodiscus procer* (BARRANDE in PERNER, 1903)

procer, *Cyrtodiscus*: (BARRANDE in PERNER, 1903)
1997c Horný: Text-figs 1b, c, p. 224
OUB: Letná F.; Praha – Vysočany
L 31991
T = *Cyrtodiscus procer* (BARRANDE in PERNER, 1903)

proxima, *Archinacella*: PERNER, 1903
1903 Perner: Pl. 1, figs 46, 47; Text-figs 9a, b, p. 36
1963a Horný: Pl. 8, figs 6-8 (*Archinacellina modesta*)
OUK: Králův Dvůr F.; Lejškov (= Chodouň)
■L 5867, holotype by monotypy; by error designated as a paratype of *Archinacella modesta* by Horný 1963a
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)

pusillus, *Temnodiscus*: BARRANDE in PERNER, 1903
1903 Perner: Pl. 86, fig. 18-21
1963b Horný: Pl. 30, fig. 2 [*Tropidodiscus (Peruniscus) pusillus*]
OML: Šárka F.; Osek
■L 5837, lectotype (SD Horný 1963b) + several paralectotypes
G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Temnodiscus*: BARRANDE in PERNER, 1903
1903 Perner: Pl. 86, figs 19, 20
1963b Horný: Pl. 30, fig. 3 [*Tropidodiscus (Peruniscus) pusillus*]
OML: Šárka F.; Osek
■L 5837, lectotype (SD Horný 1963b)
G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Temnodiscus*: BARRANDE in PERNER, 1903
1903 Perner: Pl. 86, fig. 21
OML: Šárka F.; Osek
L 5837, paralectotype
G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Temnodiscus*: BARRANDE in PERNER, 1903
1903 Perner: AD Pl. 86, figs 19-21
OML: Šárka F.; Osek
L 5841, paralectotype
G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Temnodiscus*: BARRANDE in PERNER, 1903
1903 Perner: AD Pl. 86, figs 19-21
OML: Šárka F.; Osek
L 5842, paralectotype
G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Tropidodiscus (Peruniscus)*: (BARRANDE in PERNER, 1903)
1963b Horný: Pl. 30, fig. 1
OML: Šárka F.; Osek

L 5840

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Tropidodiscus (Peruniscus)*: (BARRANDE in PERNER, 1903)

1963b Horný: Pl. 30, figs 4-7
OML: Šárka F.; Kařez – Pětidomky

L 5838

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Tropidodiscus (Peruniscus)*: (BARRANDE in PERNER, 1903)

1963b Horný: Pl. 31, fig. 1
OML: Šárka F.; Praha – Šárka

L 5839

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903) (Horný 1997g)

pusillus, *Tropidodiscus*: (BARRANDE in PERNER, 1903)

1999a Horný: Text-figs 2d, e, p. 127
OML: Šárka F.; Osek

L 32738 (non 32638)

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903)

pusillus, *Tropidodiscus*: (BARRANDE in PERNER, 1903)

1999a Horný: Text-fig. 3a, p. 128
OML: Šárka F.; Praha – Šárka

L 32737

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903)

pusillus, *Tropidodiscus*: (BARRANDE in PERNER, 1903)

1999a Horný: Text-fig. 3b, p. 128

OML: Šárka F.; Volduchy – Díly

L 32740

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903)

pusillus, *Tropidodiscus*: (BARRANDE in PERNER, 1903)

1999a Horný: Text-fig. 3c, p. 128

OML: Šárka F.; Praha – Šárka

L 32739

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903)

pusillus, *Tropidodiscus*: (BARRANDE in PERNER, 1903)

1999a Horný: Text-fig. 5b, p. 129

OMD: Dobrotivá F.; Praha – Libeň, Hercovka

L 32741

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903)

pusillus, *Tropidodiscus*: (BARRANDE in PERNER, 1903)

1999a Horný: AD Text-fig. 6, p. 129

OML: Šárka F.; Osek

L 32736

G = *Tropidodiscus pusillus* (BARRANDE in PERNER, 1903)

putzkeri, *Sinuitopsis*: HORNÝ, 1997

1997c Horný: p. 223

nomen nudum

T = *Sinuitopsis plasi* HORNÝ, 1997

reticulatus, *Sinuitites*: PERNER, 1903

1903 Perner: Pl. 56, figs 42, 43; Text-figs 37a, b; 38a, c; 39a-c, p. 65

1963b Horný: Pl. 4, figs 1, 2 (*Sinuitites reticulatus*)

OML: Šárka F.; Osek

■L 5617, lectotype (SD Horný 1963b)

G = *Sinuitites reticulatus* PERNER, 1903

reticulatus, *Sinuitites*: PERNER, 1903

1903 Perner: Pl. 86, figs 33, 34; Text-figs 40a, b, p. 66

1963b Horný: Pl. 4, fig. 5 (*Sinuites reticulatus*)

OML: Šárka F.; Osek

L 5620, paralectotype

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1903 Perner: AD Text-figs 37a, b, p. 65; AD Text-figs 38a-c, p. 65

1963b Horný: specimen mentioned on p. 74 (*Sinuites reticulatus*)

OML: Šárka F.; Osek

L 5618, paralectotype

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1963b Horný: Pl. 4, fig. 3

OML: Šárka F.; Osek

L 5621

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1963b Horný: specimen mentioned on p. 74

OML: Šárka F.; Osek

L 5619

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 3, figs 7, 8

OML: Šárka F.; Praha – Šárka, a brickyard

L 28997 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 3, fig. 9; Text-fig. 9G, p. 90

OML: Šárka F.; Osek

L 28998

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 3, figs 10, 11; Text-fig. 9A, p. 90

OML: Šárka F.; Osek

L 28999 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 4, figs 1, 2; Text-fig. 9J, p. 91

OML: Šárka F.; Praha – Šárka, a brickyard

L 29004 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 4, fig. 4; Text-fig. 6E, p. 87; Text-fig. 9I, p. 91

OML: Šárka F.; ? Osek

L 29005 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 4, fig. 5, Text-fig. 6D, p. 87; Text-fig. 9E, p. 90

OML: Šárka F.; Osek

L 29006 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 4, fig. 6; Text-fig. 9D, p. 90

OML: Šárka F.; Osek

L 29007 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 4, fig. 7; Text-fig. 9F, p. 90

OML: Šárka F.; Praha – Šárka, a brickyard

L 29008 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 5, figs 5, 6; Text-fig. 10E, p. 92

OML: Šárka F.; Osek

L 29009

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Pl. 5, figs 9, 10; Text-fig. 6A, p. 87; Text-fig. 10D, p. 92

OML: Šárka F.; Osek

L 29010

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Text-fig. 4B, p. 84

1996a Horný: Text-fig. 1B, p. 92 (*Sinuites reticulatus*)

OML: Šárka F.; Praha – Šárka, a brickyard

L 29014 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Text-fig. 6F, p. 87

OML: Šárka F.; Praha – Šárka, a brickyard

L 29016 ±

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1992 Horný: Text-fig. 10B, p. 92

OML: Šárka F.; Osek

L 29017

G = *Sinuites reticulatus* PERNER, 1903

reticulatus, Sinuites: PERNER, 1903

1996a Horný: Pl. 4, fig. 2

OML: Šárka F.; Praha – Šárka

L 30060

G = *Sinuites reticulatus* PERNER, 1903

roemeri, Carinariopsis: BARRANDE in PERNER, 1903

1903 Perner: Pl. 85, figs 14, 15

1963b Horný: Pl. 28, fig. 10 (*Bucanopsis roemeri*)

1997a Horný: p. 7 (? *Bucanopsis calypso*)

OUN: Letná F.; Trubská

L 5746, paralectotype

G = *Bucanopsis roemeri* (BARRANDE in PERNER, 1903) (Horný herein)

roemeri, Carinariopsis: BARRANDE in PERNER, 1903

1903 Perner: Pl. 88, fig. 45; Text-figs 63a, b, p. 92

1963b Horný: Pl. 29, figs 1, 2 (*Bucanopsis roemeri*)

1997a Horný: p. 7 (? *Bucanopsis calypso*)

OUN: Letná F.; Trubská

■ L 5744, lectotype

G = *Bucanopsis roemeri* (BARRANDE in PERNER, 1903) (Horný herein)

roemeri, Carinariopsis: BARRANDE in PERNER, 1903

1903 Perner: Pl. 88, fig. 46; Text-figs 63c-f, p. 92; Text-fig. 64, p. 92

1963b Horný: Pl. 29, figs 3-5 (*Bucanopsis roemeri*)

1997a Horný: p. 7 (? *Bucanopsis calypso*)

OUN: Letná F.; Trubská

L 5745, paralectotype

G = *Bucanopsis roemeri* (BARRANDE in PERNER, 1903) (Horný herein)

- sardesoni, Carinariopsis:** PERNER, 1903
 1903 Perner: Pl. 85, figs 17, 18; Text-figs 67a, b p. 94
 1963b Horný: Pl. 11, fig. 1 (*Grandostoma bohemicum*)
 OUB: Záhořany F.; Praha – Štěrboholy
 ■L 5653, holotype by monotypy (non paratype of *Grandostoma bohemicum*, error Horný 1963b)
G = Grandostoma bohemicum (PERNER, 1903)
- sculptus, Bellerophon** (s. l.) (*Cymbularia*?): BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, figs 3, 4
 1963b Horný: p. 145 (*species dubia*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
 ■L 10853, holotype by monotypy
G = ? Sinuites cf. S. aff. bilobatus (SOWERBY, 1839) (Horný herein)
- simaki, Cyrtodiscus:** HORNÝ, 1997
 1997c Horný: Textfig. 2, p. 225
 OUB: Záhořany F.; Praha – Spořilov
 ■L 31993 ±, holotype
T = Cyrtodiscus simaki HORNÝ 1997
- simaki, Cyrtodiscus:** HORNÝ, 1997
 1997c Horný: p. 225
 OUB: Záhořany F.; Praha – Spořilov
 L 31994 ±, paratype
T = Cyrtodiscus simaki HORNÝ, 1997
- simaki, Cyrtodiscus:** HORNÝ, 1997
 1997c Horný: p. 225
 OUB: Záhořany F.; Praha – Spořilov
 L 31995, paratype
T = Cyrtodiscus simaki HORNÝ, 1997
- Sinuites** sp.
 2002 Horný and Vonka: Pl. 1, figs 5, 6
 OUB: Bohdalec F.; Praha – Radotín
 L 32752
G = Sinuites sp.
- Sinuites** sp.
 2002 Horný and Vonka: Pl. 1, fig. 7
 OUB: Bohdalec F.; Praha – Radotín
 L 32753
G = Sinuites sp.
- Sinuites** sp.
 2002 Horný and Vonka: Pl. 1, fig. 8
 OUB: Bohdalec F.; Praha – Radotín
 L 32754
G = Sinuites sp.
- Sinuites** sp.
 2002 Horný and Vonka: Pl. 1, fig. 9
 OUB: Bohdalec F.; Praha – Radotín
 L 32755
G = Sinuites sp.
- Sinuites** sp.
 2002 Horný and Vonka: a specimen mentioned in the text
 OUB: Bohdalec F.; Praha – Radotín
 L 32756
G = Sinuites sp.
- Sinuites?** sp.
 1997c Horný: Text-fig. 11, p. 230
 OML/OMD: Šárka or Dobrotivá Fs, Praha – Šárka
 L 32001
G = Sinuites sp.
- Sinuitopsis** sp.
 1997c Horný: Text-figs 4a, b, p. 226
 OMD: Dobrotivá F.; Praha – Šárka
 L 31977
T = Sinuitopsis sp.
- Sinuitopsis?** sp.
 1903 Perner: Pl. 88, fig. 34
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr, Kosov
 L 10864
G = Sinuites sp. (Horný herein)
- Sinuitopsis?** sp.
 1991b Horný: Pl. 10, fig. 6; Pl. 11, figs 1-4; Text-fig. 5/10, p. 87;
 Text-fig. 10, p. 90
 O: stratum and locality unknown
 L 28640
T = Sinuitopsis sp. (Horný herein)
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: Pl. 86, figs 27-29; Text-fig. 25, p. 56; Text-figs 29a, b, p. 61
 OML: Šárka F.; Osek
 L 10843, paralectotype
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: Pl. 86, fig. 30
 OML: Šárka F.; Osek
 L 10844, paralectotype
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: Pl. 86, fig 31 (non 32)
 OML: Šárka F.; Osek
 L 10850, paralectotype
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: Pl. 86, fig. 32
 OML: Šárka F.; Osek
 L 10845, paralectotype
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: AD Pl. 86, fig. 32
 OML: Šárka F.; Osek
 L 10846, paralectotype
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: AD Pl. 86, fig. 32
 OML: Šárka F.; Osek
 L 10847, paralectotype
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: Text-figs 28a, b, p. 61
 1963b Horný: Pl. 2, figs 1-3 (*Sinuites sowerbyi*)
 OML: Šárka F.; Osek
 ■L 5610, lectotype (SD Horný 1963b)
G = Sinuites sowerbyi PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1903 Perner: Text-figs 30a, b, p. 62
 OML: Šárka F.; Osek
 L 23772, paralectotype
G = Sinuites sowerbyi PERNER, 1903

- sowerbyi, Sinuites*: PERNER, 1903
 1903 Perner: Text-fig. 31 (left), p. 62
 OML: Šárka F.; Osek
 L 23773, paralectotype
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1903 Perner: Text-fig. 31 (right), p. 62
 1963b Horný: Pl. 3, fig 6 (*Sinuites sowerbyi*)
 OML: Šárka F.; Osek
 L 5611, paralectotype
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1903 Perner: Text-figs 35a, b, p. 63
 1963b Horný: Pl. 3, fig. 2 (*Sinuites sowerbyi*)
 OML: Šárka F.; Osek
 L 5612, paralectotype
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1903 Perner: Text-figs 36a, b (non c), p. 63
 1963b Horný: Pl. 2, fig. 4; Pl. 4, fig. 6 (*Sinuites sowerbyi*)
 OML: Šárka F.; Osek
 L 5616, paralectotype
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1963b Horný: Pl. 3, fig. 1
 OML: Šárka F.; Osek
 L 5613
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1963b Horný: Pl. 3, fig. 5
 OML: Šárka F.; Praha – Šárka
 L 5614
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1963b Horný: Pl. 3, fig. 7
 OML: Šárka F.; Praha – Vokovice
 L 5615
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 1, figs 1-3, Text-fig. 5C, p. 85
 OML: Šárka F.; Praha – Šárka
 L 28985 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 1, fig. 4; Text-fig. 8D, p. 88
 OML: Šárka F.; Osek
 L 28986
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 1, figs 5, 6; Text-fig. 5E, p. 85; Text-fig. 8B, p. 88
 OML: Šárka F.; Praha – Šárka, a field
 L 28987
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 1, figs 7, 8; Text-fig. 5F, p. 85; Text-fig. 8J, p. 89
 OML: Šárka F.; Osek
 L 28988
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 2, fig. 1; Text-fig. 8I, p. 89
 OML: Šárka F.; Praha – Šárka, a brickyard
 L 28989 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 2, fig. 2; Text-fig. 8K, p. 89
 OML: Šárka F.; ? Osek
 L 28990 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 2, fig. 3; Text-fig. 8C, p. 88
 OML: Šárka F.; Osek
 L 28991
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 2, fig. 4; Text-fig. 8E, p. 88
 OML: Šárka F.; Praha – Šárka, a brickyard
 L 28992 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 2, figs 8, 9
 OML: Šárka F.; Praha – Šárka, a brickyard
 L 28993 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 2, figs 10-13; Text-figs 12A-D
 OML: Šárka F.; Praha – Šárka, a brickyard
 L 28994
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 3, figs 1-4; Text-fig. 5H, p. 85; Text-fig. 8H, p. 89
 OML: Šárka F.; Praha – Šárka, a brickyard
 L 28995 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Pl. 3, figs 5, 6; Text-fig. 3, p. 83, Text-fig. 5G, p. 85;
 Text-fig. 8L, p. 89
 OML: Šárka F.; Praha – Vokovice, a brickyard
 L 28996
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Text-fig. 1, p. 81
 OML: Šárka F.; Praha – Šárka, a field
 L 29011 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Text-fig. 2, p. 82
 OML: Šárka F.; Praha – Šárka
 L 29012 ±
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites*: PERNER, 1903
 1992 Horný: Text-fig. 4A, p. 84
 1996a Horný: Text-fig. 1A, p. 92 (*Sinuites sowerbyi*)
 OML: Šárka F.; Praha – Šárka, a brickyard
 L 29013 ±
G = *Sinuites sowerbyi* PERNER, 1903

- sowerbyi, Sinuites:** PERNER, 1903
 1992 Horný: Text-fig. 5A, p. 85; Text-fig. 10A, p. 92
 OML: Šárka F.; Praha – Šárka
L 29015
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1992 Horný: Text-fig. 8A, p. 88
 OML: Šárka F.; Praha – Šárka, a field
L 29019
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, Sinuites:** PERNER, 1903
 1996a Horný: Pl. 8, figs 1-3
 OML: Šárka F.; Volduchy
L 30061
G = *Sinuites sowerbyi* PERNER, 1903
- sowerbyi, cf., Sinuites:** PERNER, 1903
 1992 Horný: Text-fig. 10C, p. 92
 OML: Šárka F.; Osek
L 29018
G = *Sinuites cf. sowerbyi* PERNER, 1903
- sowerbyi var. crenata, Sinuites:** PERNER, 1903
 1903 Perner: Text-figs 33a, b, p. 63
 1963b Horný: p. 73 (*Sinuitopsis sowerbyi*)
 OML: Šárka F.; Osek
■L 23775, holotype by monotypy
G = *Sinuites sowerbyi* PERNER, 1903 (Horný 1963b)
- sowerbyi var. evoluta, Sinuites:** PERNER, 1903
 1903 Perner: Text-figs 34a-c, p. 63
 1963b Horný: Pl. 6, fig. 4 (*Sinuitopsis neglecta*)
 1997c Horný: p. 226 (*Sinuitopsis?*)
 OML: Šárka F.; Osek
■L 5629, holotype by monotypy
G = *Sinuitopsis evoluta* (PERNER, 1903) (Horný herein)
- strangulatus, Bellerophon (s. l.) (Sinuites?):** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, figs 12, 13
 1963b Horný: specimen mentioned on p. 76 (*Strangulites strangulatus*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5647, paralectotype
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903) (Horný herein)
- strangulatus, Bellerophon (s. l.) (Sinuites?):** BARRANDE in PERNER, 1903
 1903 Perner: Pl. 87, fig. 14
 1963b Horný: Pl. 5, fig. 6; Pl. 6, figs 1-3 (*Strangulites strangulatus*)
 1990 Horný: Pl. 2, fig. 6 [*Sinuites (Strangulites) strangulatus*]
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
■L 5646, lectotype (SD Horný 1963b)
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903) (Horný herein)
- strangulatus, Bellerophon (s. l.) (Sinuites?):** BARRANDE in PERNER, 1903
 1963b Horný: Pl. 5, figs 4, 5 (*Strangulites strangulatus*)
 1990 Horný: specimen mentioned on p. 115 [*Sinuites (Strangulites) strangulatus*]
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5650
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903) (Horný herein)
- strangulatus, Strangulites:** (BARRANDE in PERNER, 1903)
 1963b Horný: specimen mentioned on p. 76
- OUK: Králův Dvůr F.; Beroun – Králův Dvůr
L 5648
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903) (Horný herein)
- strangulatus, Sinuites (Strangulites):** (BARRANDE in PERNER, 1903)
 1990 Horný: Pl. 1, figs 1-4; Pl. 2, figs 1-4; Text-fig 4, p. 116, Textfig. 5, p. 117
 OUK: Králův Dvůr F.; Levín, Levín Hill
L 28508
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903)
- strangulatus, Sinuites (Strangulites):** (BARRANDE in PERNER, 1903)
 1990 Horný: Pl. 2, fig. 5
 OUK: Králův Dvůr F.; Levín, Levín Hill
L 28540
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903)
- strangulatus, Sinuites (Strangulites):** (BARRANDE in PERNER, 1903)
 1991a Horný: Pl. 1, figs 1-4; Text-fig. 1A
 OUB: Králův Dvůr F.; Levín, Levín Hill
L 28833
G = *Sinuites strangulatus* (BARRANDE in PERNER, 1903)
- sublaevis, Archinacella:** PERNER, 1903
 1903 Perner: Pl. 1, figs 31, 32
 1963a Horný: Pl. 8, fig. 5 (not 4) (*Archinacellina modesta*)
 OUK: Králův Dvůr F.; Beroun – Králův Dvůr
■L 5892, holotype by monotypy
G = *Archinacellina modesta* (BARRANDE in PERNER, 1903)
- taconicum, Grandostoma:** MAREK, 1963
 1963 Marek: Pl. 1, figs 1, 2
 1963b Horný: Pl. 13, fig. 1 (*Grandostoma taconicum*)
 OUK: Kosov F.; Praha – Běchovice
■L 36539, holotype
G = *Grandostoma taconicum* MAREK, 1963
- tarda, Barrandicella:** (PERNER, 1903)
 1999 Peel and Horný: Text-figs 9 A-G, p. 108
 OUB: Záhořany F.; Praha – Štěrboholy
L 31965
G = *Barrandicella?* *tarda* (PERNER, 1903) (Horný herein)
- tarda, Barrandicella:** (PERNER, 1903)
 1999 Peel and Horný: Text-figs 9 A-G, p. 109
 OUB: Záhořany F.; Praha – Dubec
L 31964
G = *Barrandicella?* *tarda* (PERNER, 1903) (Horný herein)
- Tropidodiscus** sp. A
 1997c Horný: Text-fig. 18a, p. 234
 OUB: Záhořany F.; Praha – Spořilov
L 32378
G = *Tropidodiscus* sp.
- Tropidodiscus** sp. B
 1997c Horný: Textfig. 18b, p. 234
 OUK: Králův Dvůr F.; Lejškov (= Chodouň)
L 32377
G = *Tropidodiscus* sp.
- Tryblidium** sp.: ŘÍHA, 1938
 1938 Říha: Pl. 1, figs 6a, 6b (*male*)
 1957 Horný: Pl. 4, fig. 4 (Tryblidiid sp.)
 OMD: Dobrotivá F.; Praha – Vokovice

- L 30074**, paratype of *Pentalina prantli*
T = *Pentalina prantli* HORNÝ, 1961 (Horný 1963a)
- vokovicensis*, *Plectonotus*? HORNÝ, 1963
1963b Horný: Pl. 27, figs 6, 7
OMD: Dobrotivá F.; Praha – Šárka, field at the villa
■L 5835 ±, holotype
T = *Sarkanella vokovicensis* (HORNÝ, 1963) (Horný 1997c)
- vokovicensis*, *Plectonotus*? HORNÝ, 1963
1963b Horný: Pl. 27, figs 8, 9
OMD: Dobrotivá F.; Praha – Šárka, field at the villa
L 5836, paratype
T = *Sarkanella vokovicensis* (HORNÝ, 1963) (Horný 1997c)
- vokovicensis*, *Sarkanella*: (HORNÝ, 1997)
1997c Horný: Text-fig. 6a, p. 227
OMD: Dobrotivá F.; Mýto
L 32371
T = *Sarkanella vokovicensis* (HORNÝ, 1997)

List of missing or unidentified specimens

- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
1903 Perner: Pl. 85, figs 22, 23
OUK: Králův Dvůr F.; Králův Dvůr
- grande*, *Salpingostoma*: BARRANDE in PERNER, 1903
1903 Perner: Pl. 85, fig. 24
OUK: Králův Dvůr F.; Králův Dvůr

Index of species (mandatory types in bold numbers)

- atavum*, *Salpingostoma*? **83**
bicarinatus, *Temnodiscus* **83**
bilobatus, aff., *Sinuitopsis* **83**, 87, Pl. 5/5
bilobatus, cf. aff., *Sinuitopsis* (?) **96**
bilobatus var. *infaustus*, *Sinuitopsis* **83**
bohemica, *Bucanella* **83**
bohemica, *Bucaniella* **83**
bohemica, *Bucanopsina* **85**
bohemica, *Carinariopsis* **83**
bohemica, *Eobucania*? **83**, Pl. 6/8
bohemica, *Kornoutella* **83**, 84, Pl. 4/5, 6
bohemica, *Tritonophon* **83**
bohemicum, *Grandostoma* **83**, 84, 85, 86, 96, Pl. 6/11, 12
bohemicus, *Hypseloconus*? **84**
bohemicus, *Tritonophon*? **83**, Pl. 6/6, 7
bouceki, *Tropidodiscus* **84**, Pl. 5/10
calypso, *Bucanopsina* **84**, 85, Pl. 6/2-4
calypso, *Bucanopsina* (?) **95**
calypso, cf., *Bucanopsina* **85**
calypso, *Bucanopsis* **85**
capuloidea, *Archinacella* **85**
capulooides, *Archinacella* **85**
catenularia, *Carinariopsis* **85**
comata, *Bucania* **85**
comata, cf., *Bucanopsis* **85**
consobrina, *Pterotheca* **85**, 86, Pl. 6/9
delicatula, *Bucania* **86**
draboviensis, *Bellerophon* (s.l.) (*Bucania*?) **86**
elevata, *Barrandicellopsis* **86**, 92, Pl. 3/1, 2
evoluta, *Sinuitopsis* **98**, Pl. 1/2
evoluta, cf., *Sinuitopsis* **89**
evolvens, *Cyrtodiscus*? **86**, Pl. 1/15
evolvens, *Cyrtolites*? **86**
evolvens, *Temnodiscus* **86**
extenuata, *Barrandicellopsis*? **86**, Pl. 2/5, 6
extenuata, *Orthonychia* **86**
extenuata, *Orthonychia*? **86**
ferrigena, *Temnodiscus* **86**
grande, *Grandostoma* **86**, 87, Pl. 6/13
- grande*, *Salpingostoma* **86**, **87**, **99**
grande? , *Salpingostoma* **87**
hanusi, *Sinuitopsis* **87**, Pl. 5/6
holubi, *Barrandicella* **92**, Pl. 2/13, 14
hornyi, *Sinuitopsis* Pl. 1/5, 6
immigrans, *Palaeacmaea* **87**
incertus, *Temnodiscus* **87**
incertus, cf., *Temnodiscus* **87**
incertus? , *Temnodiscus* **87**
incola, *Bellerophon* (s.l.) (*Oxydiscus*?) **88**
kettneri, *Palaeacmaea* **88**
kettneri, *Pygmaeoconus* **88**, Pl. 3/12
kettneri, *Pygmaeoconus*? **88**
kloucekii, *Cymbularia* **88**, Pl. 6/10
kolebabai, *Sarkanella* **88**, Pl. 2/7
krafti, *Pygmaeoconus* **88**, Pl. 4/2, 3
latiuscula, *Palaeacmaea* **88**
latiusculus, *Pygmaeoconus*? **88**, Pl. 4/7
mergli, *Peelerophon* **88**, Pl. 2/8, 9
mitra, *Archinacella* **88**
modesta, *Archinacella* **88**, **89**
modesta? , *Archinacella* **89**
modesta, *Archinacellina* **85**, 88, 89, 94, 98, Pl. 3/9
modesta, *Archinacellina* (?) **88**
neglecta, *Sinuitopsis* **83**, 86, 87, **89**, 90, 91, 93, 99, Pl. 1/3, 4
neglecta var. *transgrediens*, *Sinuitopsis* **90**, **99**
nitidus, *Cyrtodiscus* **91**, Pl. 1/9, 10
nitidus, *Gamadiscus* **91**
nitidus, *Oxydiscus* (*Cyrtodiscus*) **91**
nodosa, *Sinuitopsis* **91**
nodosa? , *Sinuitopsis* **86**
novaki, *Sarkanella* **92**, Pl. 2/5, 6
ordovicina, *Palaeoscurria* **92**
ordovicinus, *Micropileus*? **92**, Pl. 4/4
ovata, *Archinacella* **92**
ovata, *Barrandicella* **92**, 93, Pl. 2/10-12
ovata elevata, *Archinacella*? **92**
ovata var. *complanata*, *Archinacella* **92**
ovata var. *elevata*, *Archinacella* **92**

- ovata* var. *rostrata*, *Archinacella* **92**
ovata var. *subglobosa*, *Archinacella* **92**
ovata var. *tarda*, *Archinacella* **92**
ovata var. *typica*, *Archinacella* **92**
peeli, *Tritonophon* 83, **84**, 93, Pl. 6/5
perneri, *Selesinutes* **93**, Pl. 5/7, 8
plasi, *Sinuitopsis* **93**, 94, Pl. 1/7, 8
platynotus, *Temnodiscus* **93**
podexpulicis, *Marekicella* **93**, Pl. 3/7, 8
porrecta, *Palaeacmaea?* **93**
porrectus, *Pygmaeoconus* **93**, Pl. 4/1
pragensis, *Carcassonnella* **93**, Pl. 2/1, 2
prantli, *Pentalina* **93**, 99, Pl. 1/1
primula, *Palaeacmaea* **87**, **93**
primulus, *Patelliconus* **87**, 93, Pl. 3/10, 11
procer, *Cyrtodiscus* **93**, 94, Pl. 1/11-13
procer, *Oxydiscus* (*Cyrtodiscus*) **93**, 94
proxima, *Archinacella* **94**
pusillus, *Temnodiscus* **94**
pusillus, *Tropidodiscus* **94**, Pl. 1/10; Pl. 5/9
pusillus, *Tropidodiscus* (*Peruniscus*) **94**
putzkeri, *Sinuitopsis* **94**
reticulatus, *Sinutes* **94**, 95, Pl. 4/11; Pl. 5/1-3
roemerii, *Bucanopsis* 85 (?), 95, Pl. 6/1
roemerii, *Bucanopsis* **95**
roemerii, *Carinariopsis* **95**, 99
sardesoni, *Carinariopsis* **96**
sculptus, *Bellerophon* (s.l.) (*Cymbularia?*) **96**
simaki, *Cyrtodiscus* **96**, Pl. 1/14
Sinuitopsis? **96**
sowerbyi, *Sinutes* **96**, 97, 98, 99, Pl. 4/8-10
sowerbyi, cf., *Sinutes* **98**
sowerbyi var. *crenata*, *Sinutes* **98**
sowerbyi var. *evoluta*, *Sinutes* **98**
 sp. A, *Tropidodiscus* **98**
 sp. B, *Tropidodiscus* **98**
 sp., *Bucanella* **83**
 sp., *Bucanella?* **84**
 sp., *Bucanopsina* **84**
 sp., *Bucanopsina?* **87**
 sp., *Bucanopsis* **84**
 sp., *Cymbularia* **86**
 sp., *Sinutes* **96**
 sp., *Sinutes?* **96**
 sp., *Sinutes* (?) **88**
 sp., *Sinuitopsis* **96**
 sp., *Sinuitopsis?* **96**
 sp., *Tropidodiscus* **98**
 sp., *Tryblidiid* **98**
 sp., *Tryblidium* **98**
 sp. n., *Carcassonnella* **85**
strangulatus, *Bellerophon* (s.l.) (*Sinutes?*) **98**
strangulatus, *Sinutes* **98**, Pl. 5/4
strangulatus, *Sinutes* (*Strangulites*) **98**
strangulatus, *Strangulites* **98**
sublaevis, *Archinacella* **98**
taconicum, *Grandostoma* **98**, Pl. 6/14
tarda, *Archinacella* **92**
tarda, *Archinacella* (?) **89**
tarda, *Barrandicella* **98**
tarda, *Barrandicella?* **92**, 98, Pl. 3/3, 4
tarda, *Barrandicella?* (?) **89**
vokovicensis, *Pleconotus?* **99**
vokovicensis, *Sarkanella* **99**, Pl. 2/3, 4
-
- Archinacella modesta* **88**
Archinacella modesta? **89**
Archinacella ovata **92**
Archinacella ovata var. *complanata* **92**
Archinacella ovata var. *elevata* **92**
Archinacella ovata var. *rostrata* **92**
Archinacella ovata var. *subglobosa* **92**
Archinacella ovata var. *tarda* **92**
Archinacella ovata var. *typica* **92**
Archinacella proxima **94**
Archinacella sublaevis **98**
Archinacella tarda **92**
Archinacella tarda (?) **89**
Archinacella? *ovata elevata* **92**
Archinacellina modesta 85, 88, 89, 94, 98, Pl. 3/9
Archinacellina modesta (?) **88**
Barrandicella holubi **92**, Pl. 2/13, 14
Barrandicella ovata **92**, 93, Pl. 2/10-12
Barrandicella tarda **98**
Barrandicella? *tarda* **92**, 98, Pl. 3/3, 4
Barrandicella? *tarda* (?) **89**
Barrandicellopsis elevata 86, 92, Pl. 3/1, 2
Barrandicellopsis? *extenuata* 86, Pl. 2/5, 6
Bellerophon (s.l.) (*Bucania?*) *draboviensis* **86**
Bellerophon (s.l.) (*Cymbularia?*) *sculptus* **96**
Bellerophon (s.l.) (*Oxydiscus?*) *incola* **88**
Bellerophon (s.l.) (*Sinutes?*) *strangulatus* **98**
Bucanella bohemica **83**
Bucanella sp. **83**
Bucanella? sp. **84**
Bucania comata **85**
Bucania delicatula **86**
Bucaniella bohemica **83**
Bucanopsis bohemica **85**
Bucanopsis calypso 84, 85, Pl. 6/2-4
Bucanopsis calypso (?) **95**
Bucanopsis cf. *calypso* **85**
Bucanopsis roemerii 85(?), 95, Pl. 6/1
Bucanopsis sp. **84**
Bucanopsis? sp. **87**
Bucanopsis cf. *comata* **85**
Bucanopsis calypso **85**
Bucanopsis sp. **84**
Bucanopsis roemerii **95**
Carcassonnella pragensis **93**, Pl. 2/1, 2
Carcassonnella sp. n. **85**
Carinariopsis bohemica **83**
Carinariopsis catenularia **85**
Carinariopsis roemerii **95**, 99
Carinariopsis sardesoni **96**
Cymbularia klouckei **88**, Pl. 6/10
Cymbularia sp. **86**
Cyrtodiscus nitidus **91**, Pl. 1/9, 10
Cyrtodiscus procer **93**, 94, Pl. 1/11-13
Cyrtodiscus simaki **96**, Pl. 1/14
Cyrtodiscus? *evolvens* 86, Pl. 1/15
Cyrtolites? *evolvens* **86**
Cytonellopsis elevata **86**
Eobucania? *bohemica* **83**, Pl. 6/8
Gamadiscus nitidus **91**
Grandostoma bohemicum 83, 84, 85, 86, 96, Pl. 6/11, 12
Grandostoma grande **86**, 87, Pl. 6/13
Grandostoma taconicum **98**, Pl. 6/14
Hypseloconus? *bohemicus* **84**
Kornoutella bohemica 83, 84, Pl. 4/5, 6
Marekicella podexpulicis **93**, Pl. 3/7, 8
Micropileus? *ordovicinus* **92**, Pl. 4/4
Orthonychia extenuata **86**
Orthonychia? *extenuata* **86**
Oxydiscus (*Cyrtodiscus*) *nitidus* **91**
Oxydiscus (*Cyrtodiscus*) *procer* **93**, 94
Palaeacmaea immigrans **87**
Palaeacmaea kettneri **88**
Palaeacmaea latiuscula **88**

Index of genera (mandatory types in bold numbers)

- Archinacella capuloidea* **85**
Archinacella caploides **85**
Archinacella mitra **88**

- Palaeacmaea primula* 87, 93
Palaeacmaea? porrecta 93
Palaeoscurria ordovicina 92
Patelliconus primulus 87, 93, Pl. 3/10, 11
Peleorophon mergli 88, Pl. 2/8, 9
Pentalina prantli 93, 99, Pl. 1/1
Plectronotus? vokovicensis 99
Pterotheca consobrina 85, 86, Pl. 6/9
Pygmaeoconus kraftii 88, Pl. 4/2, 3
Pygmaeoconus porrectus 93, Pl. 4/1
Pygmaeoconus kettneri 88, Pl. 3/12
Pygmaeoconus? kettneri 88
Pygmaeoconus? latiusculus 88, Pl. 4/7
Salpingostoma grande 86, 87, 99
Salpingostoma grande? 87
Salpingostoma? atavum 83
Sarkanella kolebabai 88, Pl. 2/7
Sarkanella novaki 92, Pl. 2/5, 6
Sarkanella vokovicensis 99
Selesinuites perneri 93, Pl. 5/7, 8
Sinuites aff. bilobatus 83, 87, Pl. 5/5
Sinuites cf. aff. bilobatus (?) 96
Sinuites bilobatus var. *infaustus* 83
Sinuites hanusi 87, Pl. 5/6
Sinuites reticulatus 94, 95, Pl. 4/11; Pl. 5/1-3
Sinuites sowerbyi 96, 97, 98, 99, Pl. 4/8-10
Sinuites sowerbyi var. *crenata* 98
Sinuites sowerbyi var. *evoluta* 98
Sinuites cf. sowerbyi 98
Sinuites sp. 96
Sinuites sp.? 96
Sinuites sp. (?) 88
Sinuites strangulatus 98, Pl. 5/4
Sinuites (Strangulites) strangulatus 98
Sinuitopsis plasi 93, 94, Pl. 1/7, 8
Sinuitopsis putzkeri 94
Sinuitopsis? 96
Sinuitopsis evoluta 98, Pl. 1/2
Sinuitopsis cf. evoluta 89
Sinuitopsis hornyi Pl. 1/5, 6
Sinuitopsis neglecta 83, 86, 87, 89, 90, 91, 93, 99, Pl. 1/3, 4
Sinuitopsis neglecta var. *transgrediens* 90, 99
Sinuitopsis nodosa 91
Sinuitopsis nodosa? 86
Sinuitopsis sp. 96
Strangulites strangulatus 98
Temnodiscus bicarinatus 83
Temnodiscus evolvens 86
Temnodiscus ferrigena 86
Temnodiscus incertus 87
Temnodiscus incertus? 87
Temnodiscus cf. incertus 87
Temnodiscus platynotus 93
Temnodiscus pusillus 94
Tritonophon bohemica 83
Tritonophon peeli 83, 84, 93, Pl. 6/5
Tritonophon? *boemicus* 8/3, Pl. 6/6, 7
Tropidodiscus bouceki 84, Pl. 5/10
Tropidodiscus (Peruniscus) pusillus 94
Tropidodiscus pusillus 94, Pl. 1/10; Pl. 5/9
Tropidodiscus sp. 98
Tropidodiscus sp. A 98
Tropidodiscus sp. B 98
Tryblidiid sp. 98
Tryblidium sp. 98
- Dzik, J. (1981): Larval development, musculature, and relationships of *Sinuitopsis* and related Baltic bellerophonts. – Norsk geologisk Tidsskrift, 61: 111-121.
Ebbestad, J. O. R. (1999): Evolution and ecology of Baltic late Ordovician univalved molluscs (Comprehensive summary of Doctoral dissertation). – Uppsala University, Department of Earth Sciences, Historical Geology and Palaeontology. Uppsala.
Frýda, J. (1988): A new species of *Modestospira* (Gastropoda) from the Ordovician of Bohemia. – Věstník Ústředního ústavu geologického, 63: 227-232.
Geyer, G. (1994): Middle Cambrian mollusks from Idaho and early conchiferan evolution. – In: Landing, E. (ed.), Studies in stratigraphy and paleontology in honor of Donald W. Fisher. – New York State Museum Bulletin 481: 69-86.
Havlíček, V. (1989): Climate changes and development of benthic communities through the Mediterranean Ordovician. – Sborník geologických věd, Geologie, 44: 79-116.
Havlíček, V. (1998): Ordovician. – In: Chlupáč, I. et al., Palaeozoic of the Barrandian. V, Prague Basin. 41-79.
Havlíček, V., Vaněk, J. (1966): The biostratigraphy of the Ordovician. – Sborník geologických věd, Paleontologie, 8: 7-69.
Havlíček, V., Vaněk, J., and Fatka, O. (1994): Perunica microcontinent in the Ordovician (its position within the Mediterranean Province, series division, benthic and pelagic associations). – Sborník geologických věd, Geologie, 46: 23-56.
Holub, K. (1911): Nová fauna spodního siluru v okolí Rokycan. – Rozpravy II. třídy České akademie, 20, 15: 1-19.
Holub, K. (1912): Doplňky ku fauně Eulomového horizontu v okolí Rokycan. – Rozpravy II. třídy České akademie, 21, 33: 1-12.
Horný, R. J. (1956): Tryblidiinae PILSBRY, 1899 (Gastropoda) from the Silurian of Central Bohemia. – Sborník Ústředního ústavu geologického, oddíl paleontologický, 22: 73-102.
Horný, R. J. (1961): New genera of Bohemian Monoplacophora and Patelid Gastropoda. – Věstník Ústředního ústavu geologického, 36: 299-302.
Horný, R. J. (1962): New genera of Lower Paleozoic Bellerophontina. – Věstník Ústředního ústavu geologického, 37: 463-466.
Horný, R. J. (1963a): Lower Paleozoic Monoplacophora and Patelid Gastropoda (Mollusca) of Bohemia. – Sborník Ústředního ústavu geologického, 28(1961), oddíl paleontologický, 7-83.
Horný, R. J. (1963b): Lower Paleozoic Bellerophontina (Gastropoda) of Bohemia. – Ústřední ústav geologický, Sborník geologických věd, Paleontologie, 2: 57-164.
Horný, R. J. (1963c): O systematickém postavení cyrtonelloidních měkkýšů (Mollusca) [On the systematic position of the cyrtonelloids (Mollusca)]. – Časopis Národního muzea, oddíl přírodovědný, 132, 132, 91-96.
Horný, R. J. (1964): Nové rody gastropodů z českého staršího paleozoika (Mollusca) [New Lower Paleozoic gastropod genera of Bohemia (Mollusca)]. – Časopis Národního muzea, oddíl přírodovědný, 133, 211-216.
Horný, R. J. (1965): *Cyrtolites* CONRAD, 1838 and its position among the Monoplacophora (Molusca). – Acta Musei Nationalis Pragae, B, Historia naturalis, 21: 57-70.
Horný, R. J. (1990): Muscle scars in *Sinuites* (*Strangulites*) from the Ordovician of Bohemia. – Časopis Národního muzea v Praze, řada přírodovědná, 155(1986): 109-118.
Horný, R. J. (1991a): Additional information about the muscle scars of *Sinuites* (*Strangulites*) (Mollusca). – Časopis Národního muzea v Praze, řada přírodovědná, 156(1987): 46-48.
Horný, R. J. (1991b): Shell morphology and muscle scars of *Sinuitopsis neglecta* PERNER (Mollusca, Monoplacophora). – Časopis Národního muzea v Praze, řada přírodovědná, 157(1988): 81-105.
Horný, R. J. (1991c): Vývoj názorů na klasifikaci cyklomyárních měkkýšů (Mollusca, Monoplacophora). – Časopis Národního Muzea, řada přírodovědná, 157(3-4): 13-32.
Horný, R. J. (1992): Svalové vtisky u rodu *Sinuites* (Mollusca, Gastropoda) z českého spodního ordoviku [Muscle scars in *Sinuites*

References

Barrande, J. (1872): Système silurien du centre de la Bohême, 1, Trilobites, Supplément, texte et planches 1-35. Prague, Paris.

- (Mollusca, Gastropoda) from the Lower Ordovician of Bohemia]. – Časopis Národního muzea v Praze, řada přírodovědná, 158(1988): 79-100.
- Horný, R. J. (1996a): Secondary shell deposits and presumed mode of life in *Sinuites* (Mollusca, Gastropoda). – Acta Musei Nationalis Pragae, Series B, Historia Naturalis, 51(1995): 89-103.
- Horný, R. J. (1996b): Retractor muscle scars in *Gamadiscus* (Mollusca, Tergomya). – Věstník Českého geologického ústavu, 71: 243-247.
- Horný, R. J. (1996c): Apical muscle attachment areas found in *Archinacella ovata* BARRANDE in PERNER, 1903 (Mollusca) – a preliminary report. – Věstník Českého geologického ústavu, 71: 367-368.
- Horný, R. J. (1996d): *Grandostoma*: an additional bellerophontiform mollusc with circumbilical retractor muscle attachment area (Gastropoda, Bellerophontoidea). – Journal of the Czech Geological Society, 41: 223-231.
- Horný, R. J. (1997a): *Bucanopsis* gen. n., a new bellerophontoidan gastropod with circumbilical muscle attachment areas from the Middle Ordovician of Bohemia. – Věstník Českého geologického ústavu, 72: 5-13.
- Horný, R. J. (1997b): Shell breakage and repair in explanate bellerophontidean gastropods from the Middle Ordovician of Bohemia. – Věstník Českého geologického ústavu, 72: 157-168.
- Horný, R. J. (1997c): New, rare, and better recognized Ordovician Tergomya and Gastropoda (Mollusca) of Bohemia. – Věstník Českého geologického ústavu, 73(3): 223-237.
- Horný, R. J. (1997d): Shell breakage and repair in *Sinuitopsis neglecta* (Mollusca, Tergomya) from the Middle Ordovician of Bohemia. – Časopis Národního muzea, řada přírodovědná, 166: 137-142.
- Horný, R. J. (1997e): Circumbilical retractor muscle attachment areas in the Upper Ordovician bellerophontidean gastropod *Grandostoma grande* (BARRANDE in PERNER, 1903). – Časopis Národního muzea, řada přírodovědná, 166: 99-103.
- Horný, R. J. (1997f): Circumbilical retractor muscle attachment area in the Ordovician trilobed bellerophontidean gastropod *Tritonophon peeli* sp. n. – Věstník Českého geologického ústavu, 73(4): 333-338.
- Horný, R. J. (1997g): Ordovician Tergomya and Gastropoda (Mollusca) of the Anti-Atlas (Morocco). – Acta Musei Nationalis Pragae, Series B, Natural History, 53(3-4): 37-80.
- Horný, R. J. 1998: Fossilized intestinal contents in the Lower Ordovician *Cyrtodiscus nitidus* (Mollusca, Tergomya) from the Barrandian Area (Bohemia). – Věstník Českého geologického ústavu, 73: 211-216.
- Horný, R. J. (1999a): Circumbilical retractor muscle attachment areas found in *Tropidodiscus* (Gastropoda, Bellerophontoidea). – Journal of the Czech Geological Society, 44: 126-130.
- Horný, R. J. (1999b): Gastropod, paragastropod, and tergomyan biostratigraphy of the Ordovician of Bohemia. – Acta Universitatis Carolinae – Geologica 1999, 43 (1-2): 397-400.
- Horný, R. J. (2000a): *Barrandicelopsis*, a new Ordovician archinacelloid genus (Mollusca, Gastropoda). – Věstník Českého geologického ústavu, 75: 145-52.
- Horný, R. J. (2000b): *Barrandicella holubi* sp. n. (Gastropoda) from the Sárka Formation (Ordovician, Llanvirnian) of Bohemia. – Časopis Národního muzea, řada přírodovědná, 169: 69-73.
- Horný, R. J. (2002a): The youngest representative of *Sarkanella* (Mollusca, Tergomya) from the Králův Dvůr Formation (Upper Ordovician, Kralodvorian) of the Barrandian Area (Czech Republic). – Časopis Národního muzea, řada přírodovědná, 170; in press.
- Horný, R. J. (2002b): Ordovician Tergomya and isostrophic Gastropoda of Bohemia: Types and referred specimens in the collections of the National Museum, Prague, Czech Republic. – Acta Musei Nationale Pragae, Serie B, Historia naturalis, 57 (3-4), in press (this volume).
- Horný, R. J., Peel, J. S. (1996): *Carcassonnella*, a new Lower Ordovician bellerophontiform mollusc with dorsally located retractor muscle attachments (Class Tergomya). – Věstník Českého geologického ústavu, 71: 305-331.
- Horný, R., Vonka, V. (2002): *Sinuites* Community in the Upper Ordovician Bohdalec Formation at Praha-Radotín (Barrandian Area, Czech Republic). – Časopis Národního muzea, řada přírodovědná, 170, in press.
- Knight, J. B. (1941): Paleozoic Gastropod Genotypes. – Geological Society of America, Special Papers, 32: 1-510. Baltimore.
- Knight, J. B. (1947): Bellerophont muscle scars. – Journal of Paleontology, 21: 264-267.
- Marek, L. (1963): Bellerophontid gastropods in the Kosov Beds (Ordovician) of Bohemia. – Věstník Ústředního ústavu geologického, 38: 277-280. Praha.
- Peel, J. S. (1972): Observations on some Lower Palaeozoic tremanitiform Bellerophontacea (Gastropoda) from North America. – Palaeontology, 15: 412-422.
- Peel, J. S. (1974): Systematics, ontogeny and functional morphology of Silurian trilobed bellerophontacean gastropods. – Bulletin of the Geological Society of Denmark, 23: 231-263.
- Peel, J. S. (1976): Musculature and systematic position of *Megalomphala taenia* (Bellerophontacea, Gastropoda) from the Silurian of Gotland. – Bulletin of the Geological Society of Denmark, 25: 49-55.
- Peel, J. S. (1978): Faunal succession and mode of life of Silurian gastropods in the Arisaig group, Nova Scotia. – Palaeontology, 21: 285-306.
- Peel, J. S. (1982): Muscle scars in *Bellerophon recticostatus* (Mollusca) from the Carboniferous of Ireland. – Journal of Paleontology, 56: 1307-1310.
- Peel, J. S. (1991): The Classes Tergomya and Helcionelloida, and early molluscan evolution. – Bulletin of the Grønlands geologiske Undersøgelse, 161: 11-165.
- Peel, J. S. (1993): Muscle scars and mode of life of *Carinaropsis* (Bellerophontoidea, Gastropoda) from the Silurian of Tennessee. – Journal of Paleontology, 67: 528-534.
- Peel, J. S., Horný, R. J. (1999): Muscle scars and systematic position of the Lower Palaeozoic limpets *Archinacella* and *Barrandicella* gen. n. (Mollusca). – Journal of the Czech Geological Society, 44: 97-115.
- Perner, J. (1903, 1907, 1911): Gastéropodes. – In: Barrande, J., Système Silurien du centre de la Bohême, 4(1): I-XI, I-164; (2): I-XI, I-380; (3): I-XVII, I-390. Prague.
- Pillet, J. (1992): Les Gastéropodes de l'Ordovicien moyen en Anjou. – Bulletin de la Société des Études Scientifiques du Anjou, 14: 1-22.
- Ríha, A. (1938): Příspěvek k poznání nových plžů ze středočeského paleozoika. – Věstník Královské české společnosti nauk, Tř matematicko-přírodovědecká, 1938: 1-12.
- Wahlman, G. P. (1992): Middle and Upper Ordovician symmetrical univalved mollusks (Monoplacophora and Bellerophontina) of the Cincinnati arch region. – U. S. Geological Survey, Professional Paper, 1166-O: O1-O213. Washington.
- Webers, G. F., Pojeta, J., Jr., and Yochelson, E. L. (1992): Cambrian mollusca from the Minaret Formation, Ellsworth Mountains, West Antarctica. – In: Webers, G. F., Craddock, C., and Splettstoesser, J. F., Geology and Paleontology of the Ellsworth Mountains, West Antarctica. – Geological Society of America, Memoir 170, Chapter 10, pp. 181-248. Boulder, Colorado.
- Wenz, W. (1940): Ursprung und frühe Stammesgeschichte der Gastropoden. – Archiv der Molluskenkunde, 72: 1-10.
- Yochelson, E. L. (1982): Mollusques Monoplacophores and Gastropodes. – In: Babin, C. et al., Brachiopodes (Articulés) et Mollusques (Bivalves, Rostroconches, Monoplacophores, Gastropodes) de l'Ordovicien inférieur (Tremadocien-Arenigien) de la Montagne Noire (France méridionale). – Mémoire de la Société des Études Scientifiques de l'Aude, 1982: 51-59. Carcassonne.
- Yochelson, E. L. (1988): A new genus of Patellacea (Gastropoda) from the Middle Ordovician of Utah: The oldest known example of the superfamily. – New Mexico Bureau of Mines and Mineral Resources Memoir 44: 195-200.